



V2V EDTECH LLP

Online Coaching at an Affordable Price.

OUR SERVICES:

- Diploma in All Branches, All Subjects
- Degree in All Branches, All Subjects
- BSCIT / CS
- Professional Courses



+91 93260 50669



v2vedtech.com



V2V EdTech LLP



v2vedtech



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

Important Instructions to Examiners:

- 1) The answers should be examined by key words and not as word-to-word as given in the model answer scheme.
- 2) The model answer and the answer written by candidate may vary but the examiner may try to assess the understanding level of the candidate.
- 3) The language errors such as grammatical, spelling errors should not be given more importance (Not applicable for subject English and Communication Skills).
- 4) While assessing figures, examiner may give credit for principal components indicated in the figure. The figures drawn by candidate and model answer may vary. The examiner may give credit for any equivalent figure drawn.
- 5) Credits may be given step wise for numerical problems. In some cases, the assumed constant values may vary and there may be some difference in the candidate's answer and model answer.
- 6) In case of some questions credit may be given by judgement on part of examiner of relevant answer based on candidate's understanding.
- 7) For programming language papers, credit may be given to any other program based on equivalent concept.

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|---|
| 1. | | Attempt any <u>FIVE</u> of the following: | 10 |
| | a) | State whether the function $f(x) = \frac{e^x + e^{-x}}{2}$ is odd or even. | 02 |
| | Ans | $f(x) = \frac{e^x + e^{-x}}{2}$ $\therefore f(-x) = \frac{e^{-x} + e^{-(-x)}}{2}$ $= \frac{e^{-x} + e^x}{2}$ $= f(x)$ <p>\therefore function is even.</p> | <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> |
| | b) | If $f(x) = \frac{x^2 + 1}{x^3 - 1}$ find $f\left(\frac{1}{2}\right)$ | 02 |
| | Ans | $f(x) = \frac{x^2 + 1}{x^3 - 1}$ $\therefore f\left(\frac{1}{2}\right) = \frac{\left(\frac{1}{2}\right)^2 + 1}{\left(\frac{1}{2}\right)^3 - 1}$ $= \frac{-10}{7} \quad \text{OR} \quad -1.429$ | <p>1</p> <p>1</p> |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|---|---|-------------------------|
| 1. | c) | Find $\frac{dy}{dx}$, if $y = (x^2 + 1)^5$ | 02 |
| | Ans | $y = (x^2 + 1)^5$ $\therefore \frac{dy}{dx} = 5(x^2 + 1)^4 \cdot \frac{d}{dx}(x^2 + 1)$ $= 5(x^2 + 1)^4 \cdot (2x)$ $= 10x(x^2 + 1)^4$ | 1 1 |
| | d) | Evaluate $\int (\tan x + \cot x)^2 dx$ | 02 |
| | Ans | $\int (\tan x + \cot x)^2 dx$ $= \int (\tan^2 x + 2 \tan x \cot x + \cot^2 x) dx$ $= \int (\tan^2 x + 2 + \cot^2 x) dx$ $= \int [(\sec^2 x - 1) + 2 + (\operatorname{cosec}^2 x - 1)] dx$ $= \int (\sec^2 x + \operatorname{cosec}^2 x) dx$ $= \tan x - \cot x + c$ | 1/2 1/2 1/2 + 1/2 |
| e) | Evaluate $\int \log x dx$ | 02 | |
| Ans | $\int \log x dx = \int \log x \cdot 1 dx$ $= \log x \int 1 dx - \int \left(\int 1 dx \frac{d}{dx} \log x \right) dx$ $= \log x(x) - \int x \frac{1}{x} dx$ $= x \log x - \int 1 dx$ $= x \log x - x + c$ $= x(\log x - 1) + c$ | 1/2 1/2 1/2 | |
| f) | Find the area between the lines $y = 3x$, x -axis and the ordinates $x = 1$ and $x = 5$ | 02 | |
| Ans | $\text{Area } A = \int_a^b y dx$ $= \int_1^5 3x dx$ | 1/2 | |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme | | | | | | | | | | | | |
|--------|-----------|--|----------------------------------|---|---------------------|------|----|---|------|------|------|---|-------|------|--|
| 1. | f) | $= 3 \int_1^5 x dx$ $= 3 \left[\frac{x^2}{2} \right]_1^5$ $= 3 \left[\frac{5^2}{2} - \frac{1^2}{2} \right]$ $= 36$ | <p>1/2</p> <p>1/2</p> <p>1/2</p> | | | | | | | | | | | | |
| | g) | <p>Show that there exist a root of the equation $x^2 - 2x - 1 = 0$ in $(-1, 0)$ and find approximate value of the root by using Bisection method. (Use two iterations)</p> | 02 | | | | | | | | | | | | |
| | Ans | $x^2 - 2x - 1 = 0$ $f(x) = x^2 - 2x - 1$ $f(-1) = 2$ $f(0) = -1$ <p>root is in $(-1, 0)$</p> $\therefore x_1 = \frac{-1+0}{2} = -0.5$ $\therefore f(-0.5) = 0.25$ <p>\therefore root is in $(-0.5, 0)$</p> $\therefore x_2 = \frac{-0.5+0}{2} = -0.25$ <p style="text-align: center;">OR</p> $x^2 - 2x - 1 = 0$ $f(x) = x^2 - 2x - 1$ $f(-1) = 2$ $f(0) = -1$ <p>root is in $(-1, 0)$</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>a</th> <th>b</th> <th>$x = \frac{a+b}{2}$</th> <th>f(x)</th> </tr> </thead> <tbody> <tr> <td>-1</td> <td>0</td> <td>-0.5</td> <td>0.25</td> </tr> <tr> <td>-0.5</td> <td>0</td> <td>-0.25</td> <td>----</td> </tr> </tbody> </table> | a | b | $x = \frac{a+b}{2}$ | f(x) | -1 | 0 | -0.5 | 0.25 | -0.5 | 0 | -0.25 | ---- | <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p> |
| a | b | $x = \frac{a+b}{2}$ | f(x) | | | | | | | | | | | | |
| -1 | 0 | -0.5 | 0.25 | | | | | | | | | | | | |
| -0.5 | 0 | -0.25 | ---- | | | | | | | | | | | | |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|-----------|-----------|---|---------------------|
| 2. | | Attempt any <u>THREE</u> of the following: | 12 |
| | a) | Find $\frac{dy}{dx}$ if $\cos(x^2 + y^2) = \log(xy)$ | 04 |
| | Ans | $\cos(x^2 + y^2) = \log(xy)$ $\therefore -\sin(x^2 + y^2) \left(2x + 2y \frac{dy}{dx} \right) = \frac{1}{xy} \left(x \frac{dy}{dx} + y \right)$ $\therefore -2x \sin(x^2 + y^2) - 2y \sin(x^2 + y^2) \frac{dy}{dx} = \frac{1}{y} \frac{dy}{dx} + \frac{1}{x}$ $\therefore \frac{dy}{dx} \left(-2y \sin(x^2 + y^2) - \frac{1}{y} \right) = \frac{1}{x} + 2x \sin(x^2 + y^2)$ $\therefore \frac{dy}{dx} = \frac{\frac{1}{x} + 2x \sin(x^2 + y^2)}{-2y \sin(x^2 + y^2) - \frac{1}{y}}$ | 2 1 1 |
| | b) | If $x = a \cos^3 \theta$, $y = a \sin^3 \theta$, find $\frac{dy}{dx}$ at $\theta = \frac{\pi}{4}$ | 04 |
| | Ans | <div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> $x = a \cos^3 \theta$ $\therefore \frac{dx}{d\theta} = -3a \cos^2 \theta \sin \theta$ $\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}}$ $\frac{dy}{dx} = \frac{3a \sin^2 \theta \cos \theta}{-3a \cos^2 \theta \sin \theta}$ $\frac{dy}{dx} = -\tan \theta$ at $\theta = \frac{\pi}{4}$ $\frac{dy}{dx} = -\tan \frac{\pi}{4}$ $\frac{dy}{dx} = -1$ </div> <div style="width: 45%;"> $y = a \sin^3 \theta$ $\frac{dy}{d\theta} = 3a \sin^2 \theta (\cos \theta)$ </div> </div> | 1 1 1 |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|---|
| 2. | c) | Find the maximum and minimum value of $2x^3 - 3x^2 - 36x + 10$ | 04 |
| | Ans | Let $y = 2x^3 - 3x^2 - 36x + 10$ $\therefore \frac{dy}{dx} = 6x^2 - 6x - 36$ $\therefore \frac{d^2y}{dx^2} = 12x - 6$ Consider $\frac{dy}{dx} = 0$ $6x^2 - 6x - 36 = 0$ $x^2 - x - 6 = 0$ $\therefore x = -2, x = 3$ at $x = -2$ $\frac{d^2y}{dx^2} = 12(-2) - 6 = -30 < 0$ $\therefore y$ is maximum at $x = -2$ $y_{\max} = 2(-2)^3 - 3(-2)^2 - 36(-2) + 10$ $= 54$ at $x = 3$ $\frac{d^2y}{dx^2} = 12(3) - 6 = 30 > 0$ $\therefore y$ is minimum at $x = 3$ $y_{\min} = 2(3)^3 - 3(3)^2 - 36(3) + 10$ $= -71$ | $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ |
| | d) | A beam is bent in the form of the curve $y = 2 \sin x - \sin 2x$ Find the radius of curvature of the beam at the point $x = \frac{\pi}{2}$ | 04 |
| | Ans | $y = 2 \sin x - \sin 2x$ $\therefore \frac{dy}{dx} = 2 \cos x - 2 \cos 2x$ $\therefore \frac{d^2y}{dx^2} = -2 \sin x + 4 \sin 2x$ \therefore at $x = \frac{\pi}{2}$ $\frac{dy}{dx} = 2 \cos \left(\frac{\pi}{2} \right) - 2 \cos 2 \left(\frac{\pi}{2} \right) = 2$ | $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|-------------------------------------|
| 2. | d) | $\frac{d^2y}{dx^2} = -2\sin\left(\frac{\pi}{2}\right) + 4\sin 2\left(\frac{\pi}{2}\right) = -2$ $\therefore \text{Radius of curvature is } \rho = \frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}$ $\therefore \rho = \frac{\left[1 + (2)^2\right]^{\frac{3}{2}}}{-2}$ $\therefore \rho = -5.590 \text{ or } 5.590$ | <p>½</p> <p>1</p> <p>1</p> |
| 3. | | <p>Attempt any THREE of the following:</p> | 12 |
| | a) | <p>Find the points on the curve $y = x^3 + 3x^2 - 9x + 7$ at which tangents drawn are parallel to x-axis.</p> | 04 |
| | Ans | $y = x^3 + 3x^2 - 9x + 7$ $\frac{dy}{dx} = 3x^2 + 6x - 9$ <p>\therefore tangent is parallel to x-axis</p> <p>\therefore slope of tangent = slope of x-axis</p> $\therefore \frac{dy}{dx} = 0$ $\therefore 3x^2 + 6x - 9 = 0$ $\therefore x = 1 \ ; \ x = -3$ $\therefore y = 2 \ ; \ y = 34$ <p>\therefore points are $(1, 2)$ and $(-3, 34)$</p> | <p>1</p> <p>1</p> <p>1</p> <p>1</p> |
| | b) | <p>Differentiate $\tan^{-1}\left(\frac{2x}{1-x^2}\right)$ w.r.t. $\sin^{-1}\left(\frac{2x}{1+x^2}\right)$</p> | 04 |
| | Ans | <p>Let $u = \tan^{-1}\left(\frac{2x}{1-x^2}\right)$ and $v = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$</p> <p>Put $x = \tan \theta \Rightarrow \tan^{-1} x = \theta$</p> $u = \tan^{-1}\left(\frac{2 \tan \theta}{1 - \tan^2 \theta}\right)$ $u = \tan^{-1}(\tan 2\theta)$ $u = 2\theta$ $u = 2 \tan^{-1} x$ | <p>½</p> <p>½</p> |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|--|
| 3. | b) | $\frac{du}{dx} = \frac{2}{1+x^2}$ $v = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$ $v = \sin^{-1}\left(\frac{2 \tan \theta}{1+\tan^2 \theta}\right)$ $v = \sin^{-1}(\sin 2\theta)$ $v = 2\theta$ $v = 2 \tan^{-1} x$ $\frac{dv}{dx} = \frac{2}{1+x^2}$ $\therefore \frac{du}{dv} = \frac{dx}{dv} = \frac{1+x^2}{2}$ $\frac{du}{dx} = 1$ <p style="text-align: center;">OR</p> $\text{Let } u = \tan^{-1}\left(\frac{2x}{1-x^2}\right)$ $\therefore \frac{du}{dx} = \frac{1}{1+\left(\frac{2x}{1-x^2}\right)^2} \times \left[\frac{(1-x^2)2 - 2x(-2x)}{(1-x^2)^2} \right]$ $\therefore \frac{du}{dx} = \frac{(1-x^2)^2}{(1-x^2)^2 + 4x^2} \left[\frac{2+2x^2}{(1-x^2)^2} \right]$ $\therefore \frac{du}{dx} = \frac{2+2x^2}{(1-x^2)^2 + 4x^2}$ $\therefore \frac{du}{dx} = \frac{2(1+x^2)}{(1-x^2)^2 + 4x^2}$ $\therefore \frac{du}{dx} = \frac{2(1+x^2)}{(1+x^2)^2} = \frac{2}{1+x^2}$ $\therefore v = \sin^{-1}\left(\frac{2x}{1+x^2}\right)$ | <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: **22224**

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|---|
| 3. | b) | $\therefore \frac{dv}{dx} = \frac{1}{\sqrt{1 - \left(\frac{2x}{1+x^2}\right)^2}} \times \left[\frac{(1+x^2)2 - 2x(2x)}{(1+x^2)^2} \right]$ $\therefore \frac{dv}{dx} = \frac{(1+x^2)}{\sqrt{(1+x^2)^2 - 4x^2}} \left[\frac{2-2x^2}{(1+x^2)^2} \right]$ $\therefore \frac{dv}{dx} = \frac{(2-2x^2)}{(1+x^2)\sqrt{(1+x^2)^2 - 4x^2}}$ $\therefore \frac{dv}{dx} = \frac{2(1-x^2)}{(1+x^2)(1-x^2)}$ $\therefore \frac{dv}{dx} = \frac{2}{(1+x^2)}$ $\therefore \frac{du}{dv} = \frac{dx}{dv} = \frac{2}{(1+x^2)}$ $\therefore \frac{du}{dv} = 1$ | <p>1</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> |
| | c) | <p>Find $\frac{dy}{dx}$ if $y = (\log x)^x + x^{\cos^{-1}x}$</p> <p>Ans Let $u = (\log x)^x$</p> <p>$\log u = x \log (\log x)$</p> <p>$\frac{1}{u} \frac{du}{dx} = x \frac{1}{\log x} \frac{1}{x} + \log (\log x)$</p> <p>$\therefore \frac{du}{dx} = u \left(\frac{1}{\log x} + \log (\log x) \right)$</p> <p>$\therefore \frac{du}{dx} = (\log x)^x \left[\frac{1}{\log x} + \log (\log x) \right]$</p> <p>Let $v = x^{\cos^{-1}x}$</p> <p>$\log v = \cos^{-1}x \log x$</p> | <p>04</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1/2</p> |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|---------------------------|
| 3. | c) | $\frac{1}{v} \frac{dv}{dx} = \cos^{-1} x \left(\frac{1}{x} \right) + \log x \left(\frac{-1}{\sqrt{1-x^2}} \right)$ | 1/2 |
| | | $\therefore \frac{dv}{dx} = x^{\cos^{-1} x} \left[(\cos^{-1} x) \left(\frac{1}{x} \right) - \log x \left(\frac{1}{\sqrt{1-x^2}} \right) \right]$ | 1/2 |
| | | $\therefore \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$ $\therefore \frac{dy}{dx} = (\log x)^x \left[\frac{1}{\log x} + \log(\log x) \right] + x^{\cos^{-1} x} \left[(\cos^{-1} x) \left(\frac{1}{x} \right) - \log x \left(\frac{1}{\sqrt{1-x^2}} \right) \right]$ | 1/2 |
| | d) | Evaluate: $\int \frac{\sec x \cos ecx}{\log \tan x} dx$ | 04 |
| | Ans | $\int \frac{\sec x \cos ecx}{\log \tan x} dx$ Put $\log \tan x = t$ $\therefore \frac{1}{\tan x} \sec^2 x dx = dt$ $\therefore \frac{\cos x}{\sin x} \frac{1}{\cos^2 x} dx = dt$ $\therefore \sec x \cos ecx dx = dt$ $= \int \frac{1}{t} dt$ $= \log t + c$ $= \log(\log(\tan x)) + c$ | 1 1 1/2 1 1/2 |
| 4. | | Attempt any <u>THREE</u> of the following: | 12 |
| | a) | Evaluate : $\int \frac{1}{2x^2 + 3x + 1} dx$ | 04 |
| | Ans | $\int \frac{1}{2x^2 + 3x + 1} dx$ $= \frac{1}{2} \int \frac{1}{x^2 + \frac{3}{2}x + \frac{1}{2}} dx$ | 1/2 |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|---|--|
| 4. | a) | <p>Third term = $\left(\frac{1}{2} \times \frac{3}{2}\right)^2 = \frac{9}{16}$</p> $= \frac{1}{2} \int \frac{1}{x^2 + \frac{3}{2}x + \frac{9}{16} - \frac{9}{16} + \frac{1}{2}} dx$ $= \frac{1}{2} \int \frac{1}{\left(x + \frac{3}{4}\right)^2 - \left(\frac{1}{4}\right)^2} dx$ $= \frac{1}{2} \left[\frac{1}{2\left(\frac{1}{4}\right)} \log \left(\frac{x + \frac{3}{4} - \frac{1}{4}}{x + \frac{3}{4} + \frac{1}{4}} \right) \right] + c$ $= \log \left(\frac{2x+1}{2x+2} \right) + c$ <p style="text-align: center;"><i>OR</i></p> $\int \frac{1}{2x^2 + 3x + 1} dx = \int \frac{1}{(2x+1)(x+1)} dx$ <p>Let $\frac{1}{(2x+1)(x+1)} = \frac{A}{2x+1} + \frac{B}{x+1}$</p> $1 = A(x+1) + B(2x+1)$ <p>Put $x = \frac{-1}{2}$</p> $\therefore A = 2$ <p>Put $x = -1$</p> $\therefore B = -1$ $\frac{1}{(2x+1)(x+1)} = \frac{2}{2x+1} + \frac{-1}{x+1}$ $\int \frac{1}{(2x+1)(x+1)} dx = \int \left(\frac{2}{2x+1} + \frac{-1}{x+1} \right) dx$ $= \frac{2 \log(2x+1)}{2} - \log(x+1) + c$ $= \log(2x+1) - \log(x+1) + c$ <p style="text-align: center;"><i>OR</i></p> | <p>1</p> <p>1</p> <p>1½</p> <p>½</p> <p>½</p> <p>½</p> <p>½</p> <p>1+1</p> |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: **22224**

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|---|--|
| 4. | a) | $\int \frac{1}{2x^2 + 3x + 1} dx$ $\text{Third term} = \frac{(M.T.)^2}{4(F.T.)} = \frac{9}{8}$ $= \int \frac{1}{2x^2 + 3x + \frac{9}{8} - \frac{9}{8} + 1} dx$ $= \int \frac{1}{\left(\sqrt{2x} + \frac{3}{\sqrt{8}}\right)^2 - \frac{1}{8}} dx$ $= \int \frac{1}{\left(\sqrt{2x} + \frac{3}{\sqrt{8}}\right)^2 - \left(\frac{1}{\sqrt{8}}\right)^2} dx$ $= \frac{1}{\sqrt{2}} \left[\frac{1}{2\left(\frac{1}{\sqrt{8}}\right)} \log \left(\frac{\sqrt{2x} + \frac{3}{\sqrt{8}} - \frac{1}{\sqrt{8}}}{\sqrt{2x} + \frac{3}{\sqrt{8}} + \frac{1}{\sqrt{8}}} \right) \right] + c$ $= \log \left(\frac{2x+1}{2x+2} \right) + c$ | <p>½</p> <p>1</p> <p>1</p> <p>½</p> <p>1</p> |
| | b) | <p>Evaluate : $\int \frac{dx}{a^2 \sin^2 x + b^2 \cos^2 x}$</p> <p>Ans $\int \frac{dx}{a^2 \sin^2 x + b^2 \cos^2 x}$</p> $= \int \frac{dx / \cos^2 x}{\frac{a^2 \sin^2 x + b^2 \cos^2 x}{\cos^2 x}}$ $= \int \frac{\sec^2 x dx}{a^2 \tan^2 x + b^2}$ <div style="display: flex; align-items: center; margin-left: 100px;"> <div style="border-left: 1px solid black; padding-left: 10px; margin-left: 10px;"> $\text{Put } \tan x = t$ $\therefore \sec^2 x dx = dt$ </div> </div> $= \int \frac{dt}{a^2 t^2 + b^2}$ $= \int \frac{dt}{a^2 \left(t^2 + \frac{b^2}{a^2} \right)}$ | <p>04</p> <p>½</p> <p>½</p> <p>1</p> <p>½</p> |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: **22224**

| Q. No. | Sub Q. N. | Answer | Marking Scheme | |
|--------|-----------|---|---|-----------|
| 4. | b) | $= \frac{1}{a^2} \int \frac{dt}{t^2 + \left(\frac{b}{a}\right)^2} = \frac{1}{a^2} \cdot \frac{1}{\frac{b}{a}} \tan^{-1} \left(\frac{t}{\frac{b}{a}} \right) + c$ | 1 | |
| | | $= \frac{1}{ab} \tan^{-1} \left(\frac{a \tan x}{b} \right) + c$ | ½ | |
| | | <i>OR</i> | | |
| | | $\int \frac{dx}{a^2 \sin^2 x + b^2 \cos^2 x}$ $= \int \frac{dx / \cos^2 x}{\frac{a^2 \sin^2 x + b^2 \cos^2 x}{\cos^2 x}}$ | ½ | |
| | | $= \int \frac{\sec^2 x dx}{a^2 \tan^2 x + b^2}$ | ½ | |
| | | | <u>Put $\tan x = t$</u> <u>$\therefore \sec^2 x dx = dt$</u> | |
| | | | $= \int \frac{dt}{a^2 t^2 + b^2}$ | 1 |
| | | | $= \frac{1}{b} \tan^{-1} \left(\frac{at}{b} \right) \frac{1}{a} + c$ | 1½ |
| | | | $= \frac{1}{ab} \tan^{-1} \left(\frac{a \tan x}{b} \right) + c$ | ½ |
| | | c) | Evaluate : $\int x \operatorname{cosec}^{-1} x dx$ | 04 |
| Ans | | $\int x \operatorname{cosec}^{-1} x dx$ $= \operatorname{cosec}^{-1} x \int x dx - \int \left(\int x dx \frac{d}{dx} \operatorname{cosec}^{-1} x \right) dx$ $= \operatorname{cosec}^{-1} x \cdot \frac{x^2}{2} - \int \frac{x^2}{2} \cdot \left(\frac{-1}{x\sqrt{x^2-1}} \right) \cdot dx$ $= \operatorname{cosec}^{-1} x \cdot \frac{x^2}{2} + \frac{1}{2} \int \frac{x}{\sqrt{x^2-1}} \cdot dx$ $= \operatorname{cosec}^{-1} x \cdot \frac{x^2}{2} + \frac{1}{4} \int \frac{2x}{\sqrt{x^2-1}} \cdot dx$ $= \operatorname{cosec}^{-1} x \cdot \frac{x^2}{2} + \frac{1}{4} (2\sqrt{x^2-1}) + c$ $= \operatorname{cosec}^{-1} x \cdot \frac{x^2}{2} + \frac{1}{2} (\sqrt{x^2-1}) + c$ | ½ 1 1 1 ½ | |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|---|---|
| 4. | d) | Evaluate : $\int \frac{1}{x(2-\log x)(2\log x-1)} dx$ | 04 |
| | Ans | $\int \frac{1}{x(2-\log x)(2\log x-1)} dx$ <div style="border: 1px solid black; padding: 5px; display: inline-block;"> $Put \log x = t$ $\therefore \frac{1}{x} dx = dt$ </div> $\int \frac{1}{(2-t)(2t-1)} dt$ $\frac{1}{(2-t)(2t-1)} = \frac{A}{2-t} + \frac{B}{2t-1}$ $1 = A(2t-1) + B(2-t)$ $\therefore Put \ t = 2 \ , \ A = \frac{1}{3}$ $Put \ t = \frac{1}{2} \ , \ B = \frac{2}{3}$ $\therefore \frac{1}{(2-t)(2t-1)} = \frac{\frac{1}{3}}{2-t} + \frac{\frac{2}{3}}{2t-1}$ $\int \frac{1}{(2-t)(2t-1)} dt = \int \left(\frac{\frac{1}{3}}{2-t} + \frac{\frac{2}{3}}{2t-1} \right) dt$ $= -\frac{1}{3} \log [2-t] + \frac{2}{6} \log [2t-1] + c$ $= -\frac{1}{3} \log [2-\log x] + \frac{1}{3} \log [2\log x-1] + c$ | <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> |
| | | <p style="text-align: center;">OR</p> $\int \frac{1}{x(2-\log x)(2\log x-1)} dx$ <div style="border: 1px solid black; padding: 5px; display: inline-block;"> $Put \log x = t$ $\therefore \frac{1}{x} dx = dt$ </div> $\int \frac{1}{(2-t)(2t-1)} dt$ $= \int \frac{1}{-2t^2 + 5t - 2} dt$ $= \frac{-1}{2} \int \frac{1}{t^2 - \frac{5}{2}t + 1} dt$ | <p>1/2</p> <p>1/2</p> |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|---|---|
| 4. | d) | $= \frac{-1}{2} \int \frac{1}{t^2 - \frac{5}{2}t + \frac{25}{16} - \frac{25}{16} + 1} dt$ $= \frac{-1}{2} \int \frac{1}{\left(t - \frac{5}{4}\right)^2 - \left(\frac{3}{4}\right)^2} dt$ $= \frac{-1}{2} \frac{1}{2 \frac{3}{4}} \log \left \frac{t - \frac{5}{4} - \frac{3}{4}}{t - \frac{5}{4} + \frac{3}{4}} \right + c$ $= \frac{-1}{3} \log \left \frac{t - 2}{t - \frac{1}{2}} \right + c$ $= \frac{-1}{3} \log \left \frac{\log x - 2}{\log x - \frac{1}{2}} \right + c$ | <p>½</p> <p>1</p> <p>1</p> <p>½</p> |
| | e) | <p>-----</p> <p>Evaluate: $\int_1^4 \frac{\sqrt[3]{9-x}}{\sqrt[3]{9-x} + \sqrt[3]{x+4}} dx$</p> <p>Ans $I = \int_1^4 \frac{\sqrt[3]{9-x}}{\sqrt[3]{9-x} + \sqrt[3]{x+4}} dx$ ----- (1)</p> $I = \int_1^4 \frac{\sqrt[3]{9-(5-x)}}{\sqrt[3]{9-(5-x)} + \sqrt[3]{(5-x)+4}} dx$ $\therefore I = \int_1^4 \frac{\sqrt[3]{x+4}}{\sqrt[3]{x+4} + \sqrt[3]{9-x}} dx$ ----- (2) <p>add (1) and (2), $I + I = \int_1^4 \frac{\sqrt[3]{9-x}}{\sqrt[3]{9-x} + \sqrt[3]{x+4}} dx + \int_1^4 \frac{\sqrt[3]{x+4}}{\sqrt[3]{x+4} + \sqrt[3]{9-x}} dx$</p> $\therefore 2I = \int_1^4 \frac{\sqrt[3]{9-x} + \sqrt[3]{x+4}}{\sqrt[3]{9-x} + \sqrt[3]{x+4}} dx$ $\therefore 2I = \int_1^4 1 dx$ $\therefore 2I = (x)_1^4$ $\therefore I = \frac{3}{2}$ | <p>04</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|---|----------------------------|
| 5. | | Attempt any <u>TWO</u> of the following: | 12 |
| | a) | Find the volume of the solid generated by revolving the ellipse $\frac{x^2}{9} + \frac{y^2}{4}$ about x – axis | 06 |
| | Ans | Consider $\frac{x^2}{9} + \frac{y^2}{4} = 1$ $\therefore y^2 = \frac{4}{9}(9 - x^2)$ Volume of solid $V = \pi \int_{-a}^a y^2 dx$ $V = \pi \int_{-3}^3 \frac{4}{9}(9 - x^2) dx$ $\therefore V = 2\pi \int_0^3 \frac{4}{9}(9 - x^2) dx$ $\therefore V = \frac{8\pi}{9} \left[9x - \frac{x^3}{3} \right]_0^3$ $\therefore V = \frac{8\pi}{9} \left[\left(9(3) - \frac{3^3}{3} \right) - \left(9(0) - \frac{0^3}{3} \right) \right]$ $V = 16\pi$ (Note :If student has considered/assumed other value than 1 and attempted) (to solve the problem , give appropriate marks.) | 1 1 1 1 1 1 |
| | b) | Attempt the following: | 06 |
| | (i) | Form the diffrential equation by eliminating the arbitrary constants if $y = a \cos(\log x) + b \sin(\log x)$ | 03 |
| | Ans | $y = a \cos(\log x) + b \sin(\log x)$ $\therefore \frac{dy}{dx} = -a \sin(\log x) \frac{1}{x} + b \cos(\log x) \frac{1}{x}$ $\therefore x \frac{dy}{dx} = -a \sin(\log x) + b \cos(\log x)$ $\therefore x \frac{d^2 y}{dx^2} + \frac{dy}{dx} = -a \cos(\log x) \frac{1}{x} - b \sin(\log x) \frac{1}{x}$ $\therefore x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} = -(a \cos(\log x) + b \sin(\log x))$ | 1 1 ½ |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|--|--|----------------|
| 5. | b)(i) | $\therefore x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} = -y$ $\therefore x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + y = 0$ | 1/2 |
| | b)(ii) | Solve the differential equation: $\frac{dy}{dx} + y \tan x = \cos^2 x$ | 03 |
| | Ans | $\frac{dy}{dx} + y \tan x = \cos^2 x$ <p>Comparing with $\frac{dy}{dx} + Py = Q$</p> | 1/2 |
| | | $\therefore P = \tan x \text{ and } Q = \cos^2 x$ | 1/2 |
| | | $IF = e^{\int \tan x dx} = e^{\log(\sec x)} = \sec x$ | 1/2 |
| | | $\therefore y \cdot IF = \int Q \cdot IF dx + c$ | 1 |
| | | $y \cdot \sec x = \int \cos^2 x \sec x dx + c$ | 1 |
| | | $y \cdot \sec x = \int \cos x dx + c$ | 1 |
| | | $y \cdot \sec x = \sin x + c$ | 1 |
| | c) | In a single closed electrical circuit the current 'I' at time t is given by | 06 |
| | $E - RI - L \frac{dI}{dt} = 0.$ Find the current I at time t, given that t=0, I=0 and L,R,E are constants. | | |
| Ans | $E - RI - L \frac{dI}{dt} = 0$ | | |
| | $\therefore \frac{dI}{dt} + \frac{R}{L} I = \frac{E}{L}$ <p>Comparing with $\frac{dy}{dx} + Py = Q$</p> | 1/2 | |
| | $\therefore P = \frac{R}{L} \text{ and } Q = \frac{E}{L}$ | | |
| | $IF = e^{\int \frac{R}{L} dt} = e^{\frac{Rt}{L}}$ | 1 | |
| | $\therefore I \cdot IF = \int Q \cdot IF dt + c$ | | |
| | $I \cdot e^{\frac{Rt}{L}} = \int \frac{E}{L} e^{\frac{Rt}{L}} dt + c$ | 1 | |
| | $I \cdot e^{\frac{Rt}{L}} = \frac{E}{L} \frac{e^{\frac{Rt}{L}}}{\frac{R}{L}} + c$ | 1 | |
| | $I \cdot e^{\frac{Rt}{L}} = \frac{E}{R} e^{\frac{Rt}{L}} + c$ | | |
| | When t = 0, I = 0 | | |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|---|
| 5. | c) | $\therefore c = -\frac{E}{R}$ $I \cdot e^{\frac{R}{L}t} = \frac{E}{R} e^{\frac{R}{L}t} - \frac{E}{R}$ $I = \frac{E}{R} \left(1 - e^{-\frac{R}{L}t} \right)$ | <p>1</p> <p>½</p> <p>1</p> |
| 6. | | <p>Attempt any <u>TWO</u> of the following:</p> <p>a) Attempt the following:</p> <p>(i) Solve the following system of by equations by Jacobi's -Iteration method. (Two iterations)</p> $5x + 2y + z = 12, \quad x + 4y + 2z = 15, \quad x + 2y + 5z = 20$ <p>Ans</p> $x = \frac{1}{5}(12 - 2y - z)$ $y = \frac{1}{4}(15 - x - 2z)$ $z = \frac{1}{5}(20 - x - 2y)$ <p>Starting with $x_0 = y_0 = z_0 = 0$</p> $x_1 = 2.4$ $y_1 = 3.75$ $z_1 = 4$ $x_2 = 0.1$ $y_2 = 1.15$ $z_2 = 2.02$ | <p>12</p> <p>06</p> <p>03</p> <p>1</p> <p>1</p> |
| | a(ii) | <p>Solve the following system of equation by using Gauss-Seidel method. (Two iterations)</p> $15x + 2y + z = 18, \quad 2x + 20y - 3z = 19, \quad 3x - 6y + 25z = 22$ <p>Ans</p> $x = \frac{1}{15}(18 - 2y - z)$ $y = \frac{1}{20}(19 - 2x + 3z)$ $z = \frac{1}{25}(22 - 3x + 6y)$ | <p>03</p> <p>1</p> |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|---|---|
| 6. | c) | Find the approximate root of the equation $x^4 - x - 10 = 0$, by Newton-Raphson method (Carry out four iterations) | |
| | Ans | Let $f(x) = x^4 - x - 10$ $f(1) = -10 < 0$ $f(2) = 4 > 0$ $f'(x) = 4x^3 - 1$ Initial root $x_0 = 2$ $\therefore f'(2) = 31$ $x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} = 2 - \frac{f(2)}{f'(2)} = 1.871$ $x_2 = 1.871 - \frac{f(1.871)}{f'(1.871)} = 1.856$ $x_3 = 1.856 - \frac{f(1.856)}{f'(1.856)} = 1.856$ $x_4 = 1.856 - \frac{f(1.856)}{f'(1.856)} = 1.856$ | 1 1 1 1 |
| | | <i>OR</i> | |
| | | Let $f(x) = x^4 - x - 10$ $f(1) = -10 < 0$ $f(2) = 4 > 0$ $f'(x) = 4x^3 - 1$ Initial root $x_0 = 2$ $\therefore f'(2) = 31$ $x_i = \frac{xf'(x) - [f(x)]}{f'(x)} = \frac{x(4x^3 - 1) - [x^4 - x - 10]}{4x^3 - 1}$ $= \frac{3x^4 + 10}{4x^3 - 1}$ $x_1 = 1.871$ $x_2 = 1.856$ $x_3 = 1.856$ $x_4 = 1.856$ | 1 1 2 1/2 1/2 1/2 1/2 |
| | | <i>OR</i> | |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|--|
| 6. | c) | <p>Let $f(x) = x^4 - x - 10$</p> <p>$f(-1) = -8 < 0$</p> <p>$f(-2) = 8 > 0$</p> <p>$f'(x) = 4x^3 - 1$</p> <p>Initial root $x_0 = -2$</p> <p>$\therefore f'(-2) = -33$</p> <p>$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} = -2 - \frac{f(-2)}{f'(-2)} = -1.758$</p> <p>$x_2 = -1.758 - \frac{f(-1.758)}{f'(-1.758)} = -1.700$</p> <p>$x_3 = -1.700 - \frac{f(-1.700)}{f'(-1.700)} = -1.697$</p> <p>$x_4 = -1.697 - \frac{f(-1.697)}{f'(-1.697)} = -1.697$</p> <p style="text-align: center;">OR</p> <p>Let $f(x) = x^4 - x - 10$</p> <p>$f(-1) = -8 < 0$</p> <p>$f(-2) = 8 > 0$</p> <p>$f'(x) = 4x^3 - 1$</p> <p>Initial root $x_0 = -2$</p> <p>$\therefore f'(-2) = -33$</p> <p>$x_i = \frac{xf'(x) - [f(x)]}{f'(x)} = \frac{x(4x^3 - 1) - [x^4 - x - 10]}{4x^3 - 1}$</p> <p>$= \frac{3x^4 + 10}{4x^3 - 1}$</p> <p>$x_1 = -1.758$</p> <p>$x_2 = -1.700$</p> <p>$x_3 = -1.697$</p> <p>$x_4 = -1.697$</p> <p>-----</p> | <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> |



SUMMER – 2018 EXAMINATION

Subject Name: Applied Mathematics Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|---|----------------|
| | | <p><u>Important Note</u></p> <p><i>In the solution of the question paper, wherever possible all the possible alternative methods of solution are given for the sake of convenience. Still student may follow a method other than the given herein. In such case, first see whether the method falls within the scope of the curriculum, and then only give appropriate marks in accordance with the scheme of marking.</i></p> <p>-----</p> <p>-----</p> | |



V2V EDTECH LLP

Online Coaching at an Affordable Price.

OUR SERVICES:

- Diploma in All Branches, All Subjects
- Degree in All Branches, All Subjects
- BSCIT / CS
- Professional Courses



+91 93260 50669



v2vedtech.com



V2V EdTech LLP



v2vedtech



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

Important Instructions to Examiners:

- 1) The answers should be examined by key words and not as word-to-word as given in the model answer scheme.
- 2) The model answer and the answer written by candidate may vary but the examiner may try to assess the understanding level of the candidate.
- 3) The language errors such as grammatical, spelling errors should not be given more importance (Not applicable for subject English and Communication Skills).
- 4) While assessing figures, examiner may give credit for principal components indicated in the figure. The figures drawn by candidate and model answer may vary. The examiner may give credit for any equivalent figure drawn.
- 5) Credits may be given step wise for numerical problems. In some cases, the assumed constant values may vary and there may be some difference in the candidate's answer and model answer.
- 6) In case of some questions credit may be given by judgement on part of examiner of relevant answer based on candidate's understanding.
- 7) For programming language papers, credit may be given to any other program based on equivalent concept.

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|---|--|
| 1. | | Solve any <u>FIVE</u> of following: | 10 |
| | a) | If $f(x) = x^3 - x$, find $f(1) + f(2)$ | 02 |
| | Ans | $f(x) = x^3 - x$ $\therefore f(1) = (1)^3 - (1) = 0$ $\therefore f(2) = (2)^3 - (2) = 6$ $\therefore f(1) + f(2) = 0 + 6$ $\therefore f(1) + f(2) = 6$ | $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ |
| | b) | State whether the function $f(x) = x^3 - 3x + \sin x + x \cdot \cos x$ is even or odd. | 02 |
| | Ans | $f(x) = x^3 - 3x + \sin x + x \cdot \cos x$ $\therefore f(-x) = (-x)^3 - 3(-x) + \sin(-x) + (-x) \cdot \cos(-x)$ $= -x^3 + 3x - \sin x - x \cdot \cos x$ $= -(x^3 - 3x + \sin x + x \cdot \cos x)$ $= -f(x)$ \therefore Function is odd. | $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ |
| | c) | Find $\frac{dy}{dx}$ if $y = e^{2x} \cdot \log(x+1)$ | 02 |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|--|--|----------------|
| 1. | c) | $y = e^{2x} \cdot \log(x+1)$ | |
| | Ans | $\therefore \frac{dy}{dx} = e^{2x} \cdot \frac{1}{x+1} + \log(x+1) \cdot e^{2x} \cdot 2$ $= \frac{e^{2x}}{x+1} + 2e^{2x} \log(x+1)$ | 1+1 |
| | d) | Evaluate $\int \left(e^{2x} + \frac{1}{1+x^2} \right) dx$ | 02 |
| Ans | $\int \left(e^{2x} + \frac{1}{1+x^2} \right) dx$ $= \frac{e^{2x}}{2} + \tan^{-1} x + c$ | 1+1 | |
| e) | Evaluate $\int \frac{dx}{9x^2 - 16}$ | 02 | |
| Ans | $\int \frac{dx}{9x^2 - 16} = \frac{1}{9} \int \frac{dx}{x^2 - \frac{16}{9}}$ $= \frac{1}{9} \int \frac{dx}{x^2 - \left(\frac{4}{3}\right)^2}$ $= \frac{1}{9} \cdot \frac{1}{2 \cdot \frac{4}{3}} \log \left(\frac{x - \frac{4}{3}}{x + \frac{4}{3}} \right) + c$ $= \frac{1}{24} \log \left(\frac{3x-4}{3x+4} \right) + c$ | 1/2 | |
| | OR | | |
| | | $\int \frac{dx}{9x^2 - 16} = \int \frac{dx}{(3x)^2 - (4)^2}$ $= \frac{1}{2(4)} \cdot \frac{1}{3} \log \left(\frac{3x-4}{3x+4} \right) + c$ $= \frac{1}{24} \log \left(\frac{3x-4}{3x+4} \right) + c$ | 1 |
| | | | 1 |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|---|
| 1. | f) | Find the area enclosed by the curve $y = x^3$, x -axis and the ordinates $x = 1$ and $x = 3$ | 02 |
| | Ans | $\text{Area } A = \int_a^b y dx$ $= \int_1^3 x^3 dx$ $= \left[\frac{x^4}{4} \right]_1^3$ $= \frac{(3)^4}{4} - \frac{(1)^4}{4}$ $= \frac{81}{4} - \frac{1}{4}$ $= 20$ | <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> |
| 2. | g) | Show that the root of $x^3 - 9x + 1 = 0$ lies between 2 and 3. | 02 |
| | Ans | $f(x) = x^3 - 9x + 1$ $f(2) = (2)^3 - 9(2) + 1 = -9 < 0$ $f(3) = (3)^3 - 9(3) + 1 = 1 > 0$ $\therefore \text{Root lies between 2 and 3}$ | <p>1</p> <p>1</p> |
| | | Solve any THREE of the following: | 12 |
| | a) | If $x^2 + y^2 + 2xy - y = 0$ find $\frac{dy}{dx}$ at (1, 2) | 04 |
| | Ans | $x^2 + y^2 + 2xy - y = 0$ $2x + 2y \frac{dy}{dx} + 2 \left(x \frac{dy}{dx} + y \right) - \frac{dy}{dx} = 0$ $2x + 2y \frac{dy}{dx} + 2x \frac{dy}{dx} + 2y - \frac{dy}{dx} = 0$ $2y \frac{dy}{dx} + 2x \frac{dy}{dx} - \frac{dy}{dx} = -2x - 2y$ $(2y + 2x - 1) \frac{dy}{dx} = -2x - 2y$ $\frac{dy}{dx} = \frac{-2x - 2y}{2y + 2x - 1} = \frac{-2(x + y)}{2y + 2x - 1}$ $\left(\frac{dy}{dx} \right)_{(1,2)} = \frac{-2(1+2)}{2(2)+2(1)-1} = \frac{-6}{5} \quad \text{OR} \quad -1.2$ | <p>2</p> <p>1</p> <p>1</p> |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|--|
| 2. | b) | <p>If $x = a(2\theta - \sin 2\theta)$, $y = a(1 - \cos 2\theta)$ find $\frac{dy}{dx}$ at $\theta = \frac{\pi}{4}$</p> <p>Ans</p> $x = a(2\theta - \sin 2\theta) \qquad y = a(1 - \cos 2\theta)$ $\therefore \frac{dx}{d\theta} = a(2 - 2\cos 2\theta) \qquad \frac{dy}{d\theta} = 2a \sin 2\theta$ $\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}}$ $\therefore \frac{dy}{dx} = \frac{2a \sin 2\theta}{2a(1 - \cos 2\theta)} = \frac{\sin 2\theta}{(1 - \cos 2\theta)} \quad \text{OR} \quad \frac{dy}{dx} = \frac{\sin 2\theta}{(1 - \cos 2\theta)} = \frac{2 \sin \theta \cos \theta}{2 \sin^2 \theta} = \cot \theta$ <p>at $\theta = \frac{\pi}{4}$</p> $\therefore \frac{dy}{dx} = \frac{\sin 2\left(\frac{\pi}{4}\right)}{\left(1 - \cos 2\left(\frac{\pi}{4}\right)\right)} = \frac{\sin\left(\frac{\pi}{2}\right)}{\left(1 - \cos\left(\frac{\pi}{2}\right)\right)}$ $\therefore \frac{dy}{dx} = \frac{1}{1-0} = 1 \qquad \text{OR} \qquad \frac{dy}{dx} = \cot \frac{\pi}{4} = 1$ | <p>04</p> <p>1+1</p> <p>1</p> <p>1</p> |
| | c) | <p>Find the maximum and minimum value of $y = x^3 - \frac{15}{2}x^2 + 18x$</p> <p>Ans</p> <p>Let $y = x^3 - \frac{15}{2}x^2 + 18x$</p> $\therefore \frac{dy}{dx} = 3x^2 - 15x + 18$ $\therefore \frac{d^2y}{dx^2} = 6x - 15$ <p>Consider $\frac{dy}{dx} = 0$</p> $3x^2 - 15x + 18 = 0$ $x^2 - 5x + 6 = 0$ $\therefore x = 2 \text{ or } x = 3$ <p>at $x = 2$</p> $\frac{d^2y}{dx^2} = 6(2) - 15 = -3 < 0$ <p>$\therefore y$ is maximum at $x = 2$</p> $y_{\max} = (2)^3 - \frac{15}{2}(2)^2 + 18(2) = 14$ | <p>04</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|---|---|
| 2. | c) | <p>at $x = 3$</p> $\frac{d^2y}{dx^2} = 6(3) - 15 = 3 < 0$ <p>$\therefore y$ is minimum at $x = 3$</p> $y_{\min} = (3)^3 - \frac{15}{2}(3)^2 + 18(3) = 13.5$ | <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> |
| | d) | <p>A beam is bent in the form of the curve $y = 2 \sin x - \sin 2x$.</p> <p>Find the radius of curvature of the beam at the point $x = \frac{\pi}{2}$</p> | 04 |
| | Ans | <p>$y = 2 \sin x - \sin 2x$</p> $\therefore \frac{dy}{dx} = 2 \cos x - 2 \cos 2x$ $\therefore \frac{d^2y}{dx^2} = -2 \sin x + 4 \sin 2x$ $\left(\frac{dy}{dx}\right)_{\left(x=\frac{\pi}{2}\right)} = 2 \cos\left(\frac{\pi}{2}\right) - 2 \cos 2\left(\frac{\pi}{2}\right) = 2(0) - 2(-1) = 2$ $\left(\frac{d^2y}{dx^2}\right)_{\left(x=\frac{\pi}{2}\right)} = -2 \sin\left(\frac{\pi}{2}\right) + 4 \sin 2\left(\frac{\pi}{2}\right) = -2(1) + 4(0) = -2$ $\therefore \text{Radius of curvature is } \rho = \frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}$ $\therefore \rho = \frac{\left[1 + (2)^2\right]^{\frac{3}{2}}}{-2}$ $\therefore \rho = -5.59$ $\therefore \rho = 5.59$ | <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> |
| 3. | | <p>Solve any <u>THREE</u> of the following:</p> | 12 |
| | a) | <p>Find the equation of tangent and normal to the curve $2x^2 - xy + 3y^2 = 18$ at the point $(3,1)$</p> | 04 |
| | Ans | <p>$2x^2 - xy + 3y^2 = 18$</p> $4x - \left(x \frac{dy}{dx} + y\right) + 6y \frac{dy}{dx} = 0$ | $\frac{1}{2}$ |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|--|
| 3. | a) | $4x - x \frac{dy}{dx} - y + 6y \frac{dy}{dx} = 0$ $-x \frac{dy}{dx} + 6y \frac{dy}{dx} = -4x + y$ $(-x + 6y) \frac{dy}{dx} = -4x + y$ $\frac{dy}{dx} = \frac{-4x + y}{-x + 6y}$ <p>at (3,1)</p> $\text{Slope of tangent} = \frac{dy}{dx} = \frac{-4(3)+1}{-3+6(1)} = \frac{-11}{3}$ $\text{Slope of normal} = \frac{-1}{\frac{dy}{dx}} = \frac{3}{11}$ <p>Equation of tangent</p> $y - y_1 = m(x - x_1)$ $y - 1 = \frac{-11}{3}(x - 3)$ $3y - 3 = -11x + 33$ $11x + 3y - 36 = 0$ <p>Equation of normal</p> $y - y_1 = m(x - x_1)$ $y - 1 = \frac{3}{11}(x - 3)$ $11y - 11 = 3x - 9$ $3x - 11y + 2 = 0$ | <p>½</p> <p>½</p> <p>½</p> <p>1</p> <p>1</p> |
| | b) | <p>A manufacturer can sell x items at a price of Rs. $(330 - x)$ each. The cost of producing x items is Rs. $x^2 + 10x + 12$. Determine the number of items to be sold so that the manufacturer can make the maximum profit.</p> <p>Ans</p> <p>Selling price of x items = $(330 - x)x = 330x - x^2$</p> <p>Cost price of x items = $x^2 + 10x + 12$</p> <p>Profit = Selling price - Cost price</p> <p>Let $P = (330x - x^2) - (x^2 + 10x + 12)$</p> $= 330x - x^2 - x^2 - 10x - 12$ | 04 |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|---|--|----------------------|
| 3. | b) | $P = 320x - 2x^2 - 12$ $\therefore \frac{dP}{dx} = 320 - 4x$ Put $\frac{dP}{dx} = 0$ $320 - 4x = 0$ $\therefore x = 80$ $\frac{d^2P}{dx^2} = -4 < 0$ $\therefore \text{For maximum profit manufacturer can sell 80 items.}$ | 1 1 1 1 |
| | c) | If $x^y = e^{x-y}$ then prove that $\frac{dy}{dx} = \frac{\log x}{(1 + \log x)^2}$ | 04 |
| | Ans | $x^y = e^{x-y}$ $\log x^y = \log e^{x-y}$ $y \log x = (x - y) \log e$ $y \log x = x - y$ $y \log x + y = x$ $y(\log x + 1) = x$ $y = \frac{x}{1 + \log x}$ $\frac{dy}{dx} = \frac{(1 + \log x)(1) - x \left(\frac{1}{x}\right)}{(1 + \log x)^2}$ $\frac{dy}{dx} = \frac{1 + \log x - 1}{(1 + \log x)^2}$ $\frac{dy}{dx} = \frac{\log x}{(1 + \log x)^2}$ | 1/2 1/2 1 1 |
| d) | Evaluate $\int \frac{dx}{2x + x \cdot \log x}$ | 04 | |
| Ans | $I = \int \frac{dx}{2x + x \cdot \log x}$ $= \int \frac{dx}{x(2 + \log x)}$ | 1/2 | |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|---|--|
| 3. | d) | <p>Put $2 + \log x = t$ OR Put $\log x = t$</p> $\frac{1}{x} dx = dt$ $\therefore I = \int \frac{dt}{t}$ $= \log t + c$ $= \log(2 + \log x) + c$ | <p>½</p> <p>1</p> <p>½</p> <p>1</p> <p>½</p> |
| 4. | | <p>Solve any THREE of the following:</p> | 12 |
| | a) | <p>Evaluate : $\int \frac{dx}{x^2 + 4x + 25}$</p> | 04 |
| | Ans | $I = \int \frac{dx}{x^2 + 4x + 25}$ $T.T. = \left(\frac{1}{2} \times \text{Coeff. of } x \right)^2 = \left(\frac{1}{2} \times 4 \right)^2 = 4$ $x^2 + 4x + 25 = x^2 + 4x + 4 - 4 + 25$ $= (x+2)^2 + 21 = (x+2)^2 + (\sqrt{21})^2$ $\therefore I = \int \frac{dx}{(x+2)^2 + (\sqrt{21})^2}$ $= \frac{1}{\sqrt{21}} \tan^{-1} \left(\frac{x+2}{\sqrt{21}} \right) + c$ | 1 |
| | | <p>OR $I = \int \frac{dx}{x^2 + 4x - 4 + 4 + 25}$</p> | 1 |
| | b) | <p>Evaluate $\int \frac{dx}{2 + 3 \cos 2x}$</p> | 04 |
| | Ans | $I = \int \frac{dx}{2 + 3 \cos 2x}$ <p>Put $t = \tan x$, $\cos 2x = \frac{1-t^2}{1+t^2}$ and $dx = \frac{dt}{1+t^2}$</p> $\therefore I = \int \frac{\frac{dt}{1+t^2}}{2 + 3 \left(\frac{1-t^2}{1+t^2} \right)}$ $= \int \frac{\frac{dt}{1+t^2}}{\frac{2(1+t^2) + 3(1-t^2)}{1+t^2}}$ $= \int \frac{dt}{2 + 2t^2 + 3 - 3t^2}$ | 1 |
| | | | ½ |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|--|---|------------------|
| 4. | b) | $= \int \frac{dt}{5-t^2}$ $= \int \frac{dt}{(\sqrt{5})^2 - t^2}$ $= \frac{1}{2(\sqrt{5})} \log \left(\frac{\sqrt{5}+t}{\sqrt{5}-t} \right) + c$ $= \frac{1}{2(\sqrt{5})} \log \left(\frac{\sqrt{5}+\tan x}{\sqrt{5}-\tan x} \right) + c$ | 1 1 ½ |
| | c) | Evaluate $\int x \cdot \tan^{-1} x dx$ | 04 |
| | Ans | $\int x \cdot \tan^{-1} x dx$ $= \int \tan^{-1} x \cdot x dx$ $= \tan^{-1} x \int x dx - \int \left(\int x dx \frac{d(\tan^{-1} x)}{dx} \right) dx$ $= \tan^{-1} x \left(\frac{x^2}{2} \right) - \int \left(\frac{x^2}{2} \right) \frac{1}{1+x^2} dx$ $= \frac{x^2 \tan^{-1} x}{2} - \frac{1}{2} \int \frac{x^2}{1+x^2} dx$ $= \frac{x^2 \tan^{-1} x}{2} - \frac{1}{2} \int \frac{1+x^2-1}{1+x^2} dx$ $= \frac{x^2 \tan^{-1} x}{2} - \frac{1}{2} \int \left(1 - \frac{1}{1+x^2} \right) dx$ $= \frac{x^2 \tan^{-1} x}{2} - \frac{1}{2} (x - \tan^{-1} x) + c$ | 1 1 1 1 |
| d) | Evaluate $\int \frac{x^2+1}{(x+1)(x+2)(x-3)} dx$ | 04 | |
| Ans | $\frac{x^2+1}{(x+1)(x+2)(x-3)} = \frac{A}{x+1} + \frac{B}{x+2} + \frac{C}{x-3}$ $\therefore x^2+1 = A(x+2)(x-3) + B(x+1)(x-3) + C(x+1)(x+2)$ | ½ | |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|---|
| 4. | d) | <p>Put $x = -1 \quad \therefore A = \frac{-1}{2}$</p> <p>Put $x = -2 \quad \therefore B = 1$</p> <p>Put $x = 3 \quad \therefore C = \frac{1}{2}$</p> $\therefore \int \frac{x^2 + 1}{(x+1)(x+2)(x-3)} dx = \int \frac{-\frac{1}{2}}{x+1} + \frac{1}{x+2} + \frac{\frac{1}{2}}{x-3} dx$ $\therefore \int \frac{x^2 + 1}{(x+1)(x+2)(x-3)} dx = \frac{-1}{2} \log(x+1) + \log(x+2) + \frac{1}{2} \log(x-3) + c$ | <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2} + \frac{1}{2} + \frac{1}{2}$</p> |
| | e) | <p>-----</p> <p>Evaluate $\int_0^{\pi/2} \frac{dx}{1 + \sqrt[3]{\tan x}}$</p> <p>Ans $\int_0^{\pi/2} \frac{dx}{1 + \sqrt[3]{\tan x}}$</p> $= \int_0^{\pi/2} \frac{dx}{1 + \sqrt[3]{\frac{\sin x}{\cos x}}}$ $= \int_0^{\pi/2} \frac{dx}{1 + \frac{\sqrt[3]{\sin x}}{\sqrt[3]{\cos x}}}$ $= \int_0^{\pi/2} \frac{dx}{\frac{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}}{\sqrt[3]{\cos x}}}$ $I = \int_0^{\pi/2} \frac{\sqrt[3]{\cos x}}{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}} dx \quad \text{----- (1)}$ $= \int_0^{\pi/2} \frac{\sqrt[3]{\cos\left(\frac{\pi}{2} - x\right)}}{\sqrt[3]{\cos\left(\frac{\pi}{2} - x\right)} + \sqrt[3]{\sin\left(\frac{\pi}{2} - x\right)}} dx \quad \text{-----By property}$ $I = \int_0^{\pi/2} \frac{\sqrt[3]{\sin x}}{\sqrt[3]{\sin x} + \sqrt[3]{\cos x}} dx \quad \text{----- (2)}$ <p>Add (1) and (2)</p> | <p>04</p> <p>1</p> <p>1</p> |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|--|
| 4. | e) | $I + I = \int_0^{\pi/2} \frac{\sqrt[3]{\cos x}}{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}} dx + \int_0^{\pi/2} \frac{\sqrt[3]{\sin x}}{\sqrt[3]{\sin x} + \sqrt[3]{\cos x}} dx$ $2I = \int_0^{\pi/2} \frac{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}}{\sqrt[3]{\cos x} + \sqrt[3]{\sin x}} dx$ $2I = \int_0^{\pi/2} 1 dx$ $2I = [x]_0^{\pi/2}$ $2I = \frac{\pi}{2}$ $\therefore I = \frac{\pi}{4}$ | <p>1</p> <p>1</p> |
| 5. | a) | <p>Solve any TWO of the following:</p> <p>Find the volume of the solid generated by revolving the ellipse</p> $\frac{x^2}{9} + \frac{y^2}{4} = 1 \text{ about the } x\text{-axis}$ <p>Ans</p> $\frac{x^2}{9} + \frac{y^2}{4} = 1$ $\therefore \frac{y^2}{4} = 1 - \frac{x^2}{9}$ $\therefore y^2 = \frac{4}{9}(9 - x^2)$ <p>Volume of the solid generated by revolving the ellipse about the x-axis is given by</p> $V = \pi \int_{x=-3}^{x=3} y^2 dx$ $= \pi \int_{-3}^3 \frac{4}{9}(9 - x^2) dx$ $= 2\pi \int_0^3 \frac{4}{9}(9 - x^2) dx \quad \because \text{Function is even}$ $= 2\pi \frac{4}{9} \left(9x - \frac{x^3}{3} \right)_0^3$ $= \frac{8\pi}{9} \left[\left(9(3) - \frac{(3)^3}{3} \right) - 0 \right]$ $= \frac{8\pi}{9} (27 - 9)$ $= 16\pi$ | <p>12</p> <p>06</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|---|----------------|
| 5. | b) | Solve the following. | 06 |
| | (i) | Form the differential equation by eliminating the arbitrary constants if $y^2 = 4ax$ | 03 |
| | Ans | $y^2 = 4ax \quad \text{-----(1)}$ $2y \frac{dy}{dx} = 4a \quad \text{-----(2)}$ Put (2) in (1) $\therefore y^2 = 2y \frac{dy}{dx} x$ $\therefore y = 2x \frac{dy}{dx}$ $\therefore 2x \frac{dy}{dx} - y = 0$ <hr/> | 1 |
| | (ii) | Solve $(1+x^2)dy - (1+y^2)dx = 0$ | 03 |
| | Ans | $(1+x^2)dy - (1+y^2)dx = 0$ $(1+x^2)dy = (1+y^2)dx$ $\frac{dy}{1+y^2} = \frac{dx}{1+x^2}$ $\therefore \text{Solution is,}$ $\int \frac{dy}{1+y^2} = \int \frac{dx}{1+x^2}$ $\tan^{-1} y = \tan^{-1} x + c$ <hr/> | 1 |
| | c) | A resistance of 100Ω and inductance of 0.1 henries are connected in series with a battery of 20 volts. find the current in the circuit at any instant, if the relation between L,R and E is $L \frac{di}{dt} + Ri = E$ | 06 |
| | Ans | $L \frac{di}{dt} + Ri = E$ $\therefore \frac{di}{dt} + \frac{R}{L}i = \frac{E}{L} \quad \text{Comparing with } \frac{dy}{dx} + Py = Q$ $\therefore P = \frac{R}{L} \text{ and } Q = \frac{E}{L}$ $IF = e^{\int \frac{R}{L} dt} = e^{\frac{R}{L}t}$ $\therefore \text{Solution is } i \cdot IF = \int Q \cdot IF dt + c$ $i \cdot e^{\frac{R}{L}t} = \int \frac{E}{L} e^{\frac{R}{L}t} dt + c$ | 1/2 |
| | | | 1 |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|----------------|
| 5. | c) | $i \cdot e^{\frac{R}{L}t} = \frac{E}{L} \frac{e^{\frac{R}{L}t}}{R} + c$ | 1 |
| | | $i \cdot e^{\frac{R}{L}t} = \frac{E}{R} e^{\frac{R}{L}t} + c$ <p>Initially at $t = 0, i = 0 \therefore c = \frac{-E}{R}$</p> $\therefore i \cdot e^{\frac{R}{L}t} = \frac{E}{R} e^{\frac{R}{L}t} + \left(\frac{-E}{R}\right)$ $i = \frac{E}{R} \left(1 - e^{-\frac{R}{L}t}\right)$ <p>When $R=100, L = 0.1, E= 20$</p> $i = \frac{20}{100} \left(1 - e^{-\frac{100}{0.1}t}\right)$ $i = 0.2 \left(1 - e^{-1000t}\right)$ | 1/2 |
| 6. | a) | <p>Solve any TWO of the following:</p> <p>Solve the following</p> | 12 |
| | | <p>(i) Find the approximate root of the equation $x^2 + x - 3 = 0$ in the interval (1,2) by using Bisection method(use two iterations)</p> <p>Ans $x^2 + x - 3 = 0$ $f(x) = x^2 + x - 3$ $f(1) = -1 < 0$ $f(2) = 3 > 0$ root is in (1,2) $\therefore x_1 = \frac{1+2}{2} = 1.5$ $\therefore f(1.5) = 0.75 > 0$ \therefore root is in (1,1.5) $\therefore x_2 = \frac{1+1.5}{2} = 1.25$ OR $x^2 + x - 3 = 0$ $f(x) = x^2 + x - 3$ $f(1) = -1 < 0$ $f(2) = 3 > 0$</p> | 06 03 |
| | | | 1 |
| | | | 1 |
| | | | 1 |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme | | | | | | | | | | | | | | | | |
|--------|-----------|---|---------------------|--------|---------------------|--------|-----|-----|--|--|---|---|-----|------|---|-----|------|-------|---|
| 6. | (a)(i) | <p>root is in (1, 2)</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>a</th> <th>b</th> <th>$x = \frac{a+b}{2}$</th> <th>$f(x)$</th> </tr> </thead> <tbody> <tr> <td>-ve</td> <td>+ve</td> <td></td> <td></td> </tr> <tr> <td>1</td> <td>2</td> <td>1.5</td> <td>0.75</td> </tr> <tr> <td>1</td> <td>1.5</td> <td>1.25</td> <td>-----</td> </tr> </tbody> </table> | a | b | $x = \frac{a+b}{2}$ | $f(x)$ | -ve | +ve | | | 1 | 2 | 1.5 | 0.75 | 1 | 1.5 | 1.25 | ----- | 1 |
| | a | b | $x = \frac{a+b}{2}$ | $f(x)$ | | | | | | | | | | | | | | | |
| -ve | +ve | | | | | | | | | | | | | | | | | | |
| 1 | 2 | 1.5 | 0.75 | | | | | | | | | | | | | | | | |
| 1 | 1.5 | 1.25 | ----- | | | | | | | | | | | | | | | | |
| | (ii) | <p>Solve the following system of the equations by using Gauss elimination method $x + y + z = 6$, $2x - 3y + 3z = 5$, $3x + 2y - z = 4$</p> <p>Ans</p> $\begin{aligned} x + y + z &= 6 \\ 2x - 3y + 3z &= 5 \\ 3x + 2y - z &= 4 \end{aligned}$ $\begin{aligned} 2x + 2y + 2z &= 12 && 3x + 3y + 3z = 18 \\ 2x - 3y + 3z &= 5 && \text{and} && 3x + 2y - z = 4 \\ - & && - && \\ \hline 5y - z &= 7 && && y + 4z = 14 \end{aligned}$ $\begin{aligned} 20y - 4z &= 28 \\ y + 4z &= 14 \\ + & && \\ \hline 21y &= 42 \end{aligned}$ <p>$\therefore y = 2$</p> <p>$z = 3$</p> <p>$x = 1$</p> <p>\therefore Solution is $\{1, 2, 3\}$</p> | 03 | | | | | | | | | | | | | | | | |
| | b) | <p>Solve the following system of equations by using Gauss Seidal method (use four iterations) correct upto 3 places of decimals.</p> $x + 7y - 3z = -22, 5x - 2y + 3z = 18, 2x - y + 6z = 22$ | 06 | | | | | | | | | | | | | | | | |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|----------------|-----------|---|----------------|
| 6. | b)Ans | $5x - 2y + 3z = 18, x + 7y - 3z = -22, 2x - y + 6z = 22$ | 1 |
| | | $x = \frac{1}{5}(18 + 2y - 3z)$ | |
| | | $y = \frac{1}{7}(-22 - x + 3z)$ | 1 |
| | | $z = \frac{1}{6}(22 - 2x + y)$ | |
| | | Starting with $x_0 = y_0 = z_0 = 0$ | |
| | | $x_1 = 3.6$ | |
| | | $y_1 = -3.657$ | 1 |
| | | $z_1 = 1.857$ | |
| | | $x_2 = 1.023$ | |
| | | $y_2 = -2.493$ | 1 |
| $z_2 = 2.910$ | | | |
| $x_3 = 0.857$ | | | |
| $y_3 = -2.018$ | 1 | | |
| $z_3 = 3.045$ | | | |
| $x_4 = 0.966$ | | | |
| $y_4 = -1.976$ | 1 | | |
| $z_4 = 3.015$ | | | |
| | c) | Using Newton-Raphson method find the approximate root of the equation correct upto 3 places of decimals. $x^3 - 2x - 5 = 0$ (Use four iterations) | 06 |
| | Ans | $f(x) = x^3 - 2x - 5$ | |
| | | $f(2) = -1 < 0$ | |
| | | $f(3) = 16 > 0$ | |
| | | Root is in (2,3) | 1 |
| | | $f'(x) = 3x^2 - 2$ | 1 |
| | | Initial root $x_0 = 2$ | |
| | | $\therefore f'(2) = 10$ | |



WINTER – 2018 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q. N. | Answer | Marking Scheme |
|--------|-----------|--|--|
| 6. | c) | $x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} = 2 - \frac{f(2)}{f'(2)} = 2.1$ $x_2 = 2.1 - \frac{f(2.1)}{f'(2.1)} = 2.095$ $x_3 = 2.095 - \frac{f(2.095)}{f'(2.095)} = 2.095$ $x_4 = 2.095 - \frac{f(2.095)}{f'(2.095)} = 2.095$ <p>OR</p> $f(x) = x^3 - 2x - 5$ $f(2) = -1 < 0$ $f(3) = 16 > 0$ <p>Root is in (2,3)</p> $f'(x) = 3x^2 - 2$ <p>Initial root $x_0 = 2$</p> $x_i = \frac{xf'(x) - f(x)}{f'(x)}$ $= \frac{x(3x^2 - 2) - (x^3 - 2x - 5)}{3x^2 - 2}$ $= \frac{3x^3 - 2x - x^3 + 2x + 5}{3x^2 - 2}$ $= \frac{2x^3 + 5}{3x^2 - 2}$ $x_1 = 2.1$ $x_2 = 2.095$ $x_3 = 2.095$ $x_4 = 2.095$ | <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>2</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> |
| | | <p><u>Important Note</u></p> <p><i>In the solution of the question paper, wherever possible all the possible alternative methods of solution are given for the sake of convenience. Still student may follow a method other than the given herein. In such case, first see whether the method falls within the scope of the curriculum, and then only give appropriate marks in accordance with the scheme of marking.</i></p> | |



V2V EDTECH LLP

Online Coaching at an Affordable Price.

OUR SERVICES:

- Diploma in All Branches, All Subjects
- Degree in All Branches, All Subjects
- BSCIT / CS
- Professional Courses



+91 93260 50669



v2vedtech.com



V2V EdTech LLP



v2vedtech



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

Important Instructions to Examiners:

- 1) The answers should be examined by key words and not as word-to-word as given in the model answer scheme.
- 2) The model answer and the answer written by candidate may vary but the examiner may try to assess the understanding level of the candidate.
- 3) The language errors such as grammatical, spelling errors should not be given more Importance (Not applicable for subject English and Communication Skills).
- 4) While assessing figures, examiner may give credit for principal components indicated in the figure. The figures drawn by candidate and model answer may vary. The examiner may give credit for any equivalent figure drawn.
- 5) Credits may be given step wise for numerical problems. In some cases, the assumed constant values may vary and there may be some difference in the candidate's answers and model answer.
- 6) In case of some questions credit may be given by judgement on part of examiner of relevant answer based on candidate's understanding.
- 7) For programming language papers, credit may be given to any other program based on equivalent concept.

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|--|--|-------------------------------------|
| 1. | | Solve any FIVE of the following: | 10 |
| | a) | If $f(x) = x^3 - 5x^2 - 4x + 20$ show that $f(0) = -2f(3)$ | 02 |
| | Ans | $f(x) = x^3 - 5x^2 - 4x + 20$ $\therefore f(0) = (0)^3 - 5(0)^2 - 4(0) + 20 = 20$ $\therefore f(3) = (3)^3 - 5(3)^2 - 4(3) + 20$ $= -10$ $\therefore -2f(3) = -2 \times -10 = 20 = f(0)$ | $\frac{1}{2}$ $\frac{1}{2}$ 1 |
| b) | State whether the function $f(x) = x^3 - 3x + \sin x + x \cos x$, is odd or even. | 02 | |
| Ans | $f(x) = x^3 - 3x + \sin x + x \cos x$ $\therefore f(-x) = (-x)^3 - 3(-x) + \sin(-x) + (-x)\cos(-x)$ $= -x^3 + 3x - \sin x - x \cos x$ $= -(x^3 - 3x + \sin x + x \cos x)$ $= -f(x)$ \therefore Given function is odd. | $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ | |
| c) | If $y = \sin x \cdot \cos 2x$, find $\frac{dy}{dx}$ | 02 | |
| Ans | $y = \sin x \cdot \cos 2x$ | | |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|---|---|----------------|
| 1. | c) | $\therefore \frac{dy}{dx} = \sin x(-\sin 2x) \times 2 + \cos 2x \cos x$ $= -2 \sin x \sin 2x + \cos 2x \cos x$ | 02 |
| | d) | Evaluate: $\int \cos^2 x dx$ | 02 |
| | Ans | $\int \cos^2 x dx = \int \frac{1 + \cos 2x}{2} dx$ $= \frac{1}{2} \int (1 + \cos 2x) dx$ $= \frac{1}{2} \left(x + \frac{\sin 2x}{2} \right) + c$ | 1 |
| | e) | Evaluate: $\int \frac{1}{3x+5} dx$ | 02 |
| | Ans | $\int \frac{1}{3x+5} dx$ $= \frac{1}{3} \log(3x+5) + c$ | 02 |
| f) | Find the area between the the line $y = 2x$, x -axis and ordinates $x = 1$ to $x = 3$. | 02 | |
| Ans | $\text{Area } A = \int_a^b y dx$ $= \int_1^3 2x dx$ $= 2 \left[\frac{x^2}{2} \right]_1^3 \quad \text{or} \quad [x^2]_1^3$ $= 2 \left[\frac{9}{2} - \frac{1}{2} \right] \quad \text{or} \quad [3^2 - 1^2]$ $= 8$ | 1/2 1/2 1/2 1/2 | |
| g) | Find approximate root of the equation $x^2 + x - 3 = 0$ in $(1, 2)$ by using Bisection method. (Use two iterations) | 02 | |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme | | | | | | | | | | | | | | | |
|-----------|----------|--|--|--------|---|---------------------|--------|---|---|---|-----|------|----|---|-----|------|--|---|
| 1. | g)Ans | <p>Let $f(x) = x^2 + x - 3$ $f(1) = -1$ $f(2) = 3$ \therefore the root is in $(1, 2)$ $x_1 = \frac{a+b}{2} = \frac{1+2}{2} = 1.5$ $f(1.5) = 0.75 > 0$ $x_2 = \frac{x_1+a}{2} = \frac{1.5+1}{2} = 1.25$ OR Let $f(x) = x^2 + x - 3$ $f(1) = -1, f(2) = 3 \therefore$ the root is in $(1, 2)$</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>Iteration</th> <th>a</th> <th>b</th> <th>$x = \frac{a+b}{2}$</th> <th>$f(x)$</th> </tr> </thead> <tbody> <tr> <td>I</td> <td>1</td> <td>2</td> <td>1.5</td> <td>0.75</td> </tr> <tr> <td>II</td> <td>1</td> <td>1.5</td> <td>1.25</td> <td></td> </tr> </tbody> </table> | Iteration | a | b | $x = \frac{a+b}{2}$ | $f(x)$ | I | 1 | 2 | 1.5 | 0.75 | II | 1 | 1.5 | 1.25 | | <p>$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $1\frac{1}{2}$</p> |
| Iteration | a | b | $x = \frac{a+b}{2}$ | $f(x)$ | | | | | | | | | | | | | | |
| I | 1 | 2 | 1.5 | 0.75 | | | | | | | | | | | | | | |
| II | 1 | 1.5 | 1.25 | | | | | | | | | | | | | | | |
| 2. | | <p>Solve any THREE of the following :</p> <p>a) Find $\frac{dy}{dx}$ if $x^3 + xy^2 = y^3 + yx^2$</p> <p>Ans $x^3 + xy^2 = y^3 + yx^2$ $x(x^2 + y^2) = y(y^2 + x^2)$ $x = y$ $\frac{dy}{dx} = 1$ OR $x^3 + xy^2 = y^3 + yx^2$ $3x^2 + 2xy \frac{dy}{dx} + y^2 = 3y^2 \frac{dy}{dx} + 2xy + x^2 \frac{dy}{dx}$ $\frac{dy}{dx}(2xy - 3y^2 - x^2) = 2xy - 3x^2 - y^2$</p> | <p>12 04 1 1 2 2 1</p> | | | | | | | | | | | | | | | |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q. N. | Answers | Marking Scheme |
|--------|--|---|---------------------|
| 2. | a) | $\frac{dy}{dx} = \frac{2xy - 3x^2 - y^2}{2xy - 3y^2 - x^2}$ | 1 |
| | b) | Find $\frac{dy}{dx}$ at $\theta = \frac{\pi}{4}$ if $x = a \cos^3 \theta$, $y = b \sin^3 \theta$ | 04 |
| | Ans | $x = a \cos^3 \theta$ $\therefore \frac{dx}{d\theta} = 3a \cos^2 \theta (-\sin \theta)$ $= -3a \cos^2 \theta \sin \theta$ $y = b \sin^3 \theta$ $\therefore \frac{dy}{d\theta} = 3b \sin^2 \theta \cos \theta$ $= 3b \sin^2 \theta \cos \theta$ $\therefore \frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}}$ $= \frac{3b \sin^2 \theta \cos \theta}{-3a \cos^2 \theta \sin \theta}$ $= -\frac{b}{a} \tan \theta$ $\text{at } \theta = \frac{\pi}{4}$ $\frac{dy}{dx} = -\frac{b}{a} \tan \frac{\pi}{4}$ $= -\frac{b}{a}$ | 1 1 1 |
| c) | A manufacture can sell x items per week at price $(23 - 0.001x)$ rupees each. It cost $(5x + 2000)$ rupees to produce x items Find the number items to be produced eper week for maximum profit. | 04 | |
| Ans | <p>Let number of item be x</p> <p>Selling price = $(23 - 0.001x)x$</p> $= 23x - 0.001x^2$ | $\frac{1}{2}$ | |



| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|---|--|
| 2. | c) | <p>Cost price = $(5x + 2000)$ profit = selling price – cost price $\therefore p = 23x - 0.001x^2 - (5x + 2000)$ $= 23x - 0.001x^2 - 5x - 2000$ $= 18x - 0.001x^2 - 2000$ $\frac{dp}{dx} = 18 - 0.002x$ $\frac{d^2p}{dx^2} = -0.002$ \therefore profit is maximum Let $18 - 0.002x = 0$ $x = \frac{18}{0.002} = 9000$</p> | <p>$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 1</p> |
| | d) | <p>Find the radius of curvature of the curve $y = e^x$ at the point where it crosses the Y-axis.</p> | 04 |
| | Ans | <p>$y = e^x$ $\frac{dy}{dx} = e^x$ $\frac{d^2y}{dx^2} = e^x$ curve crosses Y-axis $\therefore x = 0$ $\frac{dy}{dx} = e^0 = 1$ $\frac{d^2y}{dx^2} = e^0 = 1$ $\rho = \frac{\left(1 + \left(\frac{dy}{dx}\right)^2\right)^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}$ $= \frac{(1 + 1^2)^{\frac{3}{2}}}{1} = 2^{\frac{3}{2}} = 2.828$</p> | <p>1 1 $\frac{1}{2}$ $\frac{1}{2}$ 1</p> |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|-----------|--|---|
| 3. | b)Ans | $y = x^x + 5^x + x^5 + 5^5$ $\text{Let } u = x^x$ $\log u = \log x^x$ $\log u = x \log x$ $\frac{1}{u} \frac{du}{dx} = x \cdot \frac{1}{x} + \log x \cdot 1$ $\frac{du}{dx} = u(1 + \log x) = x^x(1 + \log x)$ $\therefore \frac{dy}{dx} = x^x(1 + \log x) + 5^x \log 5 + 5x^4$ | <p>1/2</p> <p>1</p> <p>1/2</p> <p>2</p> |
| | c) Ans | <p>If $x^3 \cdot y^2 = (x + y)^5$, show that $\frac{dy}{dx} = \frac{y}{x}$</p> $x^3 \cdot y^2 = (x + y)^5$ $\log(x^3 \cdot y^2) = \log(x + y)^5$ $\log x^3 + \log y^2 = 5 \log(x + y)$ $3 \log x + 2 \log y = 5 \log(x + y)$ $3 \frac{1}{x} + 2 \frac{1}{y} \frac{dy}{dx} = 5 \frac{1}{x + y} \left(1 + \frac{dy}{dx}\right)$ $\frac{3}{x} + \frac{2}{y} \frac{dy}{dx} = \frac{5}{x + y} + \frac{5}{x + y} \frac{dy}{dx}$ $\frac{2}{y} \frac{dy}{dx} - \frac{5}{x + y} \frac{dy}{dx} = \frac{5}{x + y} - \frac{3}{x}$ $\frac{dy}{dx} \left(\frac{2}{y} - \frac{5}{x + y} \right) = \frac{5x - 3x - 3y}{x(x + y)}$ $\frac{dy}{dx} \left(\frac{2x + 2y - 5y}{y(x + y)} \right) = \frac{5x - 3x - 3y}{x(x + y)}$ $\frac{dy}{dx} \left(\frac{2x - 3y}{y} \right) = \frac{2x - 3y}{x}$ $\frac{dy}{dx} = \frac{y}{x}$ | <p>04</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> <p>1</p> |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|--|--|
| 3. | d) | <p>Evaluate: $\int \frac{e^x(x+1)}{\sin^2(xe^x)} dx.$</p> <p>Ans $\int \frac{e^x(x+1)}{\sin^2(xe^x)} dx$</p> <p>put $xe^x = t$</p> <p>$(xe^x + e^x \cdot 1) dx = dt$</p> <p>$e^x(x+1) dx = dt$</p> <p>$= \int \frac{dt}{\sin^2 t}$</p> <p>$= \int \operatorname{cosec}^2 t dt$</p> <p>$= -\cot t + c$</p> <p>$= -\cot(xe^x) + c$</p> | <p>04</p> <p>2</p> <p>½</p> <p>1</p> <p>½</p> |
| 4 | | <p>Solve any THREE of the following:</p> <p>a) Evaluate: $\int \frac{x-3}{x^3-3x^2-16x+48} dx$</p> <p>$\int \frac{x-3}{x^3-3x^2-16x+48} dx$</p> <p>$= \int \frac{x-3}{(x-3)(x-4)(x+4)} dx$</p> <p>$= \int \frac{dx}{(x-4)(x+4)}$</p> <p>Consider $\frac{1}{(x-4)(x+4)} = \frac{A}{x-4} + \frac{B}{x+4}$</p> <p>$1 = A(x+4) + B(x-4)$</p> <p>put $x = 4$ $A = \frac{1}{8}$,</p> <p>put $x = -4$ $B = -\frac{1}{8}$</p> | <p>04</p> <p>½</p> <p>½</p> <p>½</p> |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|---|---|
| 4. | a) | $\therefore \int \frac{dx}{(x-4)(x+4)}$ $= \int \left(\frac{\frac{1}{8}}{x-4} + \frac{-\frac{1}{8}}{x+4} \right) dx$ $= \frac{1}{8} (\log(x-4) - \log(x+4)) + c$ | 2 |
| | b) | <p>Evaluate : $\int \frac{1}{2+3\cos x} dx$</p> <p>Ans $\int \frac{1}{2+3\cos x} dx$</p> <p>Put $\tan \frac{x}{2} = t$ $\cos x = \frac{1-t^2}{1+t^2}$, $dx = \frac{2dt}{1+t^2}$</p> $\therefore \int \frac{dx}{2+3\cos x} = \int \frac{1}{2+3\left(\frac{1-t^2}{1+t^2}\right)} \cdot \frac{2dt}{1+t^2}$ $= 2 \int \frac{1}{5-t^2} dt$ $= 2 \int \frac{1}{(\sqrt{5})^2 - t^2} dt$ $= 2 \times \frac{1}{2\sqrt{5}} \log \left(\frac{\sqrt{5}+t}{\sqrt{5}-t} \right) + c$ $= \frac{1}{\sqrt{5}} \log \left(\frac{\sqrt{5} + \tan \frac{x}{2}}{\sqrt{5} - \tan \frac{x}{2}} \right) + c$ | 04 1 1 ½ 1 ½ |
| | c) | <p>Evalute: $\int e^x \cdot \sin 4x dx$</p> <p>Ans $\int e^x \cdot \sin 4x dx$</p> | 04 |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|-----------|--|---|
| 4. | c) | $= \sin 4x \int e^x dx - \int \left(\int e^x dx \frac{d}{dx} \sin 4x \right) dx$ $= \sin 4x e^x - \int \cos 4x \cdot 4 \cdot e^x dx$ $= \sin 4x e^x - 4 \left[\cos 4x \int e^x dx - \int \left(\int e^x dx \cdot \frac{d}{dx} \cos 4x \right) dx \right]$ $= \sin 4x e^x - 4 \left[\cos 4x e^x - \int (-\sin 4x \cdot 4 \cdot e^x) dx \right]$ $= \sin 4x e^x - 4 \left[\cos 4x e^x + 4 \int \sin 4x \cdot e^x dx \right]$ $= \sin 4x e^x - 4 \cos 4x e^x - 16I$ $I + 16I = \sin 4x e^x - 4 \cos 4x e^x$ $17I = \sin 4x e^x - 4 \cos 4x e^x$ $I = \frac{1}{17} (\sin 4x e^x - 4 \cos 4x e^x)$ | <p>1</p> <p>½</p> <p>1</p> <p>½</p> <p>½</p> <p>½</p> |
| | d) Ans | <p>Evaluate: $\int \frac{e^x}{(e^x - 1)(e^x + 1)} dx$</p> $\int \frac{e^x}{(e^x - 1)(e^x + 1)} dx$ <p>put $e^x = t$ $e^x dx = dt$</p> $\int \frac{e^x}{(e^x - 1)(e^x + 1)} dx = \int \frac{dt}{(t-1)(t+1)}$ <p>consider $\frac{1}{(t-1)(t+1)} = \frac{A}{t-1} + \frac{B}{t+1}$</p> $1 = A(t+1) + B(t-1)$ <p>put $t = 1, A = \frac{1}{2}$</p> <p>put $t = -1, B = -\frac{1}{2}$</p> | <p>04</p> <p>1</p> <p>½</p> <p>½</p> |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|--|------------------------|
| 4. | d) | $\frac{1}{(t-1)(t+1)} = \frac{1}{2} \frac{1}{t-1} + \frac{-1}{2} \frac{1}{t+1}$ $\int \frac{dt}{(t-1)(t+1)} = \int \left(\frac{1}{2} \frac{1}{t-1} + \frac{-1}{2} \frac{1}{t+1} \right) dt$ $= \frac{1}{2} \log(t-1) - \frac{1}{2} \log(t+1) + c$ $= \frac{1}{2} \log(e^x - 1) - \frac{1}{2} \log(e^x + 1) + c$ | 1 ½ |
| e) | | <p>Evaluate : $\int_0^{\frac{\pi}{2}} \frac{1}{1 + \sqrt{\tan x}} dx$</p> <p>Ans $I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + \sqrt{\tan x}} dx$</p> $= \int_0^{\frac{\pi}{2}} \frac{1}{1 + \frac{\sqrt{\sin x}}{\sqrt{\cos x}}} dx$ $I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \text{-----(1)}$ <p>by property</p> $\therefore I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos\left(\frac{\pi}{2} - x\right)}}{\sqrt{\cos\left(\frac{\pi}{2} - x\right)} + \sqrt{\sin\left(\frac{\pi}{2} - x\right)}} dx$ $= \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \text{-----(2)}$ <p>add (1) and (2)</p> $2I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x} + \sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$ | 04 ½ 1 1 ½ |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|---|---|
| 4. | e) | $2I = \int_0^{\frac{\pi}{2}} 1 dx$ $\therefore 2I = [x]_0^{\frac{\pi}{2}}$ $= \frac{\pi}{2} - 0$ $I = \frac{\pi}{4}$ <p>OR</p> $I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + \sqrt{\tan x}} dx \text{-----(1)}$ <p>by property</p> $I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + \sqrt{\tan\left(\frac{\pi}{2} - x\right)}} dx$ $I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + \sqrt{\cot x}} dx$ $\therefore I = \int_0^{\frac{\pi}{2}} \frac{1}{1 + \sqrt{\frac{1}{\tan x}}} dx$ $= \int_0^{\frac{\pi}{2}} \frac{\sqrt{\tan x}}{\sqrt{\tan x} + 1} dx \text{-----(2)}$ <p>add (1) and (2)</p> $2I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\tan x} + 1}{\sqrt{\tan x} + 1} dx$ $2I = \int_0^{\frac{\pi}{2}} 1 dx$ $\therefore 2I = [x]_0^{\frac{\pi}{2}}$ $= \frac{\pi}{2} - 0$ $I = \frac{\pi}{4}$ | <p>½</p> <p>½</p> <p>1</p> <p>1</p> <p>½</p> <p>½</p> <p>½</p> <p>½</p> |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|---|---|----------------|
| 5. | Ans | $y = A.e^x + B.e^{-x}$ $\therefore \frac{dy}{dx} = A.e^x - B.e^{-x}$ $\therefore \frac{d^2y}{dx^2} = A.e^x + B.e^{-x}$ $\therefore \frac{d^2y}{dx^2} = y$ $\therefore \frac{d^2y}{dx^2} - y = 0$ | 1 1 1 |
| | (ii) | Solve $\frac{dy}{dx} + y \cot x = \operatorname{cosec} x$ | 03 |
| | Ans | $\frac{dy}{dx} + y \cot x = \operatorname{cosec} x$ $\therefore \text{Comparing with } \frac{dy}{dx} + Py = Q$ $P = \cot x, \quad Q = \operatorname{cosec} x$ $\text{Integrating factor } IF = e^{\int \cot x dx}$ $= e^{\log(\sin x)}$ $= \sin x$ $\therefore y \cdot IF = \int Q \cdot IF dx + c$ $\therefore y \sin x = \int \operatorname{cosec} x \cdot \sin x dx$ $\therefore y \sin x = \int 1 dx$ $\therefore y \sin x = x + c$ | |
| c) | The velocity of a particle is given by $\frac{dx}{dt} = 3t^2 - 6t + 8$. Find distance covered in 2 seconds given that $x = 0$ at $t = 0$ | | |
| | Ans | $\frac{dx}{dt} = 3t^2 - 6t + 8$ $\therefore dx = (3t^2 - 6t + 8) dt$ $\therefore \int dx = \int (3t^2 - 6t + 8) dt$ | 1 |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|-----------|--|--|
| 6. | b)Ans | $\begin{array}{rcl} 6x - y - z = 19 & & 36x - 6y - 6z = 114 \\ 3x + 4y + z = 26 & \text{and} & x + 2y + 6z = 22 \\ + \underline{\hspace{2cm}} & & + \underline{\hspace{2cm}} \\ 9x + 3y = 45 & & 37x - 4y = 136 \\ 3x + y = 15 & & 37x - 4y = 136 \\ \\ 12x + 4y = 60 & & \\ 37x - 4y = 136 & & \\ + \underline{\hspace{2cm}} & & \\ 49x = 196 & & \\ \therefore x = 4 & & \\ y = 3 & & \\ z = 2 & & \end{array}$ <p><i>Note: In the above solution, first x is eliminated and then z is eliminated to find the value of y first. If in case the problem is solved by elimination of another unknown i. e., either first y or z, appropriate marks to be given as per above scheme of marking.</i></p> | <p>1+1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> |
| | c) Ans | <p>Using Newton-Raphson method to find the approximate value of $\sqrt[3]{100}$ (perform 4 iterations)</p> <p>Let $x = \sqrt[3]{100}$</p> <p>$\therefore x^3 - 100 = 0$</p> <p>$f(x) = x^3 - 100$</p> <p>$f(4) = -36 < 0$</p> <p>$f(5) = 25 > 0$</p> <p>$f'(x) = 3x^2$</p> <p>Initial root $x_0 = 5$</p> <p>$\therefore f'(5) = 75$</p> <p>$x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} = 5 - \frac{f(5)}{f'(5)} = 4.6667$</p> <p>$x_2 = 4.6667 - \frac{f(4.6667)}{f'(4.6667)} = 4.6417$</p> | <p>06</p> <p>½</p> <p>½</p> <p>½</p> <p>½</p> <p>1</p> <p>1</p> |



SUMMER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|---|--|
| 6. | c) | $x_3 = 4.6417 - \frac{f(4.6417)}{f'(4.6417)} = 4.6416$ $x_4 = 4.6416 - \frac{f(4.6416)}{f'(4.6416)} = 4.6416$ <p>OR</p> <p>Let $f(x) = x^3 - 100$</p> <p>$f(4) = -36 < 0$ $f(5) = 25 > 0$</p> <p>$f'(x) = 3x^2$</p> <p>Initial root $x_0 = 5$</p> $x_i = x - \frac{f(x)}{f'(x)} = x - \frac{x^3 - 100}{3x^2}$ $= \frac{3x^3 - x^3 + 100}{3x^2}$ $= \frac{2x^3 + 100}{3x^2}$ <p>$x_1 = 4.6667$</p> <p>$x_2 = 4.6417$</p> <p>$x_3 = 4.6416$</p> <p>$x_4 = 4.6416$</p> <hr/> <p><u>Important Note</u></p> <p><i>In the solution of the question paper, wherever possible all the possible alternative methods of solution are given for the sake of convenience. Still student may follow a method other than the given herein. In such case, first see whether the method falls within the scope of the curriculum, and then only give appropriate marks in accordance with the scheme of marking.</i></p> <hr/> | <p>1</p> <p>1</p> <p>½</p> <p>½</p> <p>½</p> <p>½</p> <p>2</p> <p>½</p> <p>½</p> <p>½</p> <p>½</p> |



V2V EDTECH LLP

Online Coaching at an Affordable Price.

OUR SERVICES:

- Diploma in All Branches, All Subjects
- Degree in All Branches, All Subjects
- BSCIT / CS
- Professional Courses



+91 93260 50669



v2vedtech.com



V2V EdTech LLP



v2vedtech



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: **22224**

Important Instructions to Examiners:

- 1) The answers should be examined by key words and not as word-to-word as given in the model answer scheme.
- 2) The model answer and the answer written by candidate may vary but the examiner may try to assess the understanding level of the candidate.
- 3) The language errors such as grammatical, spelling errors should not be given more Importance (Not applicable for subject English and Communication Skills).
- 4) While assessing figures, examiner may give credit for principal components indicated in the figure. The figures drawn by candidate and model answer may vary. The examiner may give credit for any equivalent figure drawn.
- 5) Credits may be given step wise for numerical problems. In some cases, the assumed constant values may vary and there may be some difference in the candidate's answers and model answer.
- 6) In case of some questions credit may be given by judgement on part of examiner of relevant answer based on candidate's understanding.
- 7) For programming language papers, credit may be given to any other program based on equivalent concept.

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|--|--|
| 1. | | Solve any <u>FIVE</u> of the following: | 10 |
| | a) | State whether the function is odd or even, $f(x) = \frac{e^x + e^{-x}}{2}$ | 02 |
| | Ans | $f(x) = \frac{e^x + e^{-x}}{2}$ $\therefore f(-x) = \frac{e^{-x} + e^{-(-x)}}{2}$ $\therefore f(-x) = \frac{e^{-x} + e^x}{2}$ $\therefore f(-x) = f(x)$ $\therefore \text{function is even.}$ <p>-----</p> | ½ ½ ½ ½ |
| | b) | If $f(x) = \log_4 x + 3$, find $f\left(\frac{1}{4}\right)$ | 02 |
| | Ans | $f(x) = \log_4 x + 3$ $f\left(\frac{1}{4}\right) = \log_4\left(\frac{1}{4}\right) + 3$ $= -\log_4 4 + 3$ $= -1 + 3 = 2$ | 1 1 |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|---|--|----------------|
| 1. | c) | Find $\frac{dy}{dx}$ if $y = x^2 \cdot e^x$ | 02 |
| | Ans | $y = x^2 \cdot e^x$ $\frac{dy}{dx} = x^2 \cdot e^x + e^x \cdot 2x$ $\frac{dy}{dx} = xe^x(x+2)$ | 1 1 |
| | d) | Evaluate $\int [e^x + a^x + x^a + a^a] dx$ | 02 |
| | Ans | $\int [e^x + a^x + x^a + a^a] dx$ $= e^x + \frac{a^x}{\log a} + \frac{x^{a+1}}{a+1} + a^a x + c$ | 2 |
| e) | Evaluate: $\int \left[\frac{1}{1 + \cos 2x} \right] dx$ | 02 | |
| Ans | $\int \left[\frac{1}{1 + \cos 2x} \right] dx$ $= \int \left[\frac{1}{2 \cos^2 x} \right] dx$ $= \frac{1}{2} \int \sec^2 x dx$ $= \frac{1}{2} \tan x + c$ | 1 1 | |
| f) | Find the area bounded by $y = x$, X-axis and $x = 0$ to $x = 4$. | 02 | |
| Ans | Area $A = \int_a^b y dx$ $= \int_0^4 x dx$ $= \left[\frac{x^2}{2} \right]_0^4$ | ½ ½ | |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|--|--------------------------------|
| 1. | | $= \left(\frac{4^2}{2} - 0 \right)$ $= 8$ | $\frac{1}{2}$ $\frac{1}{2}$ |
| | g) | Find a real root of the equation $x^3 + 4x - 9 = 0$ in the interval (1, 2) by using Bisection method. (only one iteration) | 02 |
| | Ans | Let $f(x) = x^3 + 4x - 9$ $f(1) = -4$ $f(2) = 7$ \therefore the root is in (1, 2) $x_1 = \frac{a+b}{2} = \frac{1+2}{2} = 1.5$ | 1 1 |
| 2 | | Solve any THREE of the following: | 12 |
| | a) | Find $\frac{dy}{dx}$, if $y = \frac{5e^x}{3e^x + 1}$ at $x = 0$ | 04 |
| | Ans | $y = \frac{5e^x}{3e^x + 1}$ $\frac{dy}{dx} = \frac{(3e^x + 1)5e^x - 5e^x(3e^x)}{(3e^x + 1)^2}$ $\frac{dy}{dx} = \frac{15e^{2x} + 5e^x - 15e^{2x}}{(3e^x + 1)^2}$ $\frac{dy}{dx} = \frac{5e^x}{(3e^x + 1)^2}$ at $x = 0$ $\frac{dy}{dx} = \frac{5e^0}{(3e^0 + 1)^2}$ $= \frac{5}{16} \text{ or } 0.3125$ | 2 1 1 |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q. N. | Answers | Marking Scheme |
|--------|-----------|---|-------------------------------------|
| 2. | b) | If $x = a(1 + \cos \theta)$, $y = a(1 - \cos \theta)$, find $\frac{dy}{dx}$ | 04 |
| | Ans | $x = a(1 + \cos \theta)$, $y = a(1 - \cos \theta)$ $\frac{dx}{d\theta} = a(-\sin \theta) = -a \sin \theta$, $\frac{dy}{d\theta} = a(0 + \sin \theta) = a \sin \theta$ $\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}} = \frac{a \sin \theta}{-a \sin \theta}$ $\frac{dy}{dx} = -1$ | 1+1 1 1 |
| | c) | A metal wire 36 cm long is bent to form a rectangle. Find its dimensions when its area is maximum. | 04 |
| | Ans | Let length of rectangle = x , breadth = y $\therefore 2x + 2y = 36$ $\therefore y = 18 - x$ Area $A = x \times y$ $A = x(18 - x)$ $\therefore A = 18x - x^2$ $\therefore \frac{dA}{dx} = 18 - 2x$ $\therefore \frac{d^2A}{dx^2} = -2$ Let $\frac{dA}{dx} = 0$ $\therefore 18 - 2x = 0$ $\therefore x = 9$ at $x = 9$ $\frac{d^2A}{dx^2} = -2 < 0$ Area is maximum at $x = 9$ Length = 9 ; breadth = 9 | 1 1 1/2 1/2 1/2 |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|--|--|
| 2. | d) | Find radius of curvature of a curve $y = \log(\sin x)$ at $x = \pi/2$ | 04 |
| | Ans | $y = \log(\sin x)$ $\therefore \frac{dy}{dx} = \frac{1}{\sin x} \cdot \cos x = \cot x$ $\therefore \frac{d^2y}{dx^2} = -\operatorname{cosec}^2 x$ at $x = \pi/2$ $\frac{dy}{dx} = \cot \frac{\pi}{2} = 0$ $\frac{d^2y}{dx^2} = -\operatorname{cosec}^2 \frac{\pi}{2} = -1$ \therefore Radius of curvature is, $\rho = \frac{\left[1 + \left(\frac{dy}{dx}\right)^2\right]^{\frac{3}{2}}}{\frac{d^2y}{dx^2}}$ $\therefore \rho = \frac{\left[1 + (0)^2\right]^{\frac{3}{2}}}{-1}$ $\therefore \rho = -1$ i.e. 1 | $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ 1 1 |
| 3. | | Solve any <u>THREE</u> of the following: | 12 |
| | a) | Find equation of the tangent and normal to the curve $4x^2 + 9y^2 = 40$ at point (1,2) | 04 |
| | Ans | $4x^2 + 9y^2 = 40$ $\therefore 8x + 18y \frac{dy}{dx} = 0$ $\therefore \frac{dy}{dx} = \frac{-8x}{18y}$ $\therefore \frac{dy}{dx} = \frac{-4x}{9y}$ at (1,2) $\therefore \frac{dy}{dx} = \frac{-4(1)}{9(2)}$ | $\frac{1}{2}$ $\frac{1}{2}$ |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|---|--|--|
| 3. | a) | $\therefore \frac{dy}{dx} = \frac{-2}{9}$ $\therefore \text{slope of tangent, } m = \frac{-2}{9}$ <p>Equation of tangent at (1,2) is</p> $y - 2 = \frac{-2}{9}(x - 1)$ $\therefore 9y - 18 = -2x + 2$ $\therefore 2x + 9y - 20 = 0$ $\therefore \text{slope of normal, } m' = \frac{-1}{m} = \frac{9}{2}$ <p>Equation of normal at (1,2) is</p> $y - 2 = \frac{9}{2}(x - 1)$ $\therefore 2y - 4 = 9x - 9$ $\therefore 9x - 2y - 5 = 0$ | <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> |
| | b) | <p>Find $\frac{dy}{dx}$ if $y = \tan^{-1} \left[\frac{2x}{1+35x^2} \right]$</p> | 04 |
| | Ans | $y = \tan^{-1} \left[\frac{7x - 5x}{1 + 7x \cdot 5x} \right]$ $y = \tan^{-1} 7x - \tan^{-1} 5x$ $\frac{dy}{dx} = \frac{7}{1+49x^2} - \frac{5}{1+25x^2}$ | <p>1</p> <p>1</p> <p>2</p> |
| c) | <p>If $x^y = e^{x-y}$ Show that $\frac{dy}{dx} = \frac{\log x}{(1 + \log x)^2}$</p> | 04 | |
| Ans | $x^y = e^{x-y}$ $\log x^y = \log e^{x-y}$ $y \log x = x - y \log e$ $y \log x = x - y$ $y \log x + y = x$ | <p>1/2</p> <p>1/2</p> <p>1/2</p> | |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|---|-------------------------------|
| 3. | c) | $y(\log x + 1) = x$ $y = \frac{x}{\log x + 1}$ $\frac{dy}{dx} = \frac{(\log x + 1) \cdot 1 - x \cdot \frac{1}{x}}{(\log x + 1)^2}$ $= \frac{\log x}{(\log x + 1)^2}$ | 1 1 ½ |
| | d) | <p>Evaluate $\int \frac{dx}{5 + 3 \cos 2x}$</p> <p>Ans $\int \frac{dx}{5 + 3 \cos 2x}$</p> <p>Put $\tan x = t$, $dx = \frac{dt}{1+t^2}$</p> $\cos 2x = \frac{1-t^2}{1+t^2}$ $\int \frac{\frac{dt}{1+t^2}}{5 + 3 \left(\frac{1-t^2}{1+t^2} \right)}$ $= \int \frac{dt}{5(1+t^2) + 3(1-t^2)}$ $= \int \frac{dt}{5 + 5t^2 + 3 - 3t^2}$ $= \int \frac{dt}{2t^2 + 8}$ $= \int \frac{dt}{(\sqrt{2}t)^2 + (\sqrt{8})^2} \quad \text{OR} \quad = \frac{1}{2} \int \frac{dt}{t^2 + 4}$ $= \frac{1}{\sqrt{8}} \tan^{-1} \left(\frac{\sqrt{2}t}{\sqrt{8}} \right) \cdot \frac{1}{\sqrt{2}} + c \quad = \frac{1}{2} \cdot \frac{1}{2} \tan^{-1} \left(\frac{t}{2} \right) + c$ | 04 1 1 1 |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|--|--|
| 3. | d) | $= \frac{1}{\sqrt{16}} \tan^{-1} \left(\frac{\sqrt{2} \tan x}{\sqrt{8}} \right) + c \quad \text{OR} \quad = \frac{1}{4} \tan^{-1} \left(\frac{t}{2} \right) + c$ $= \frac{1}{4} \tan^{-1} \left(\frac{\tan x}{2} \right) + c \quad \text{OR} \quad = \frac{1}{4} \tan^{-1} \left(\frac{\tan x}{2} \right) + c$ | <p>1/2</p> <p>1/2</p> |
| 4. | | <p>Solve any THREE of the following:</p> | 12 |
| | a) | <p>Evaluate $\int \frac{[e^x(x+1)]}{\cos^2(x.e^x)} dx$</p> | 04 |
| | Ans | $\int \frac{[e^x(x+1)]}{\cos^2(x.e^x)} dx$ <p>Put $x.e^x = t$ $\therefore (x.e^x + e^x.1) dx = dt$ $[e^x(x+1)] dx = dt$ $\therefore \int \frac{1}{\cos^2 t} dt$ $= \int \sec^2 t dt$ $= \tan t + c$ $= \tan(x.e^x) + c$</p> | <p>1</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1/2</p> |
| | b) | <p>Evaluate: $\int \frac{dx}{2x^2 + 3x + 2}$</p> | 04 |
| | Ans | $\int \frac{dx}{2x^2 + 3x + 2}$ $= \frac{1}{2} \int \frac{dx}{x^2 + \frac{3}{2}x + 1}$ $= \frac{1}{2} \int \frac{dx}{x^2 + \frac{3}{2}x + \frac{9}{16} + 1 - \frac{9}{16}}$ | <p>1/2</p> <p>1</p> |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|---|---|
| 4. | b) | $= \frac{1}{2} \int \frac{dx}{\left(x + \frac{3}{4}\right)^2 + \frac{7}{16}}$ $= \frac{1}{2} \int \frac{dx}{\left(x + \frac{3}{4}\right)^2 + \left(\frac{\sqrt{7}}{4}\right)^2}$ $= \frac{1}{2} \cdot \frac{1}{\frac{\sqrt{7}}{4}} \tan^{-1} \left(\frac{x + \frac{3}{4}}{\frac{\sqrt{7}}{4}} \right) + c$ $= \frac{2}{\sqrt{7}} \tan^{-1} \left(\frac{4x + 3}{\sqrt{7}} \right) + c$ | <p>½</p> <p>1</p> <p>1</p> |
| | c) | <p>-----</p> <p>Evaluate $\int x^2 \cdot \tan x \, dx$</p> <p>Ans $\int x^2 \cdot \tan x \, dx$</p> $= x^2 \left(\int \tan x \, dx \right) - \int \left(\int \tan x \, dx \cdot \frac{d}{dx}(x^2) \right) dx$ $= x^2 \log(\sec x) - \int \log(\sec x) \cdot 2x \, dx$ $= x^2 \log(\sec x) - 2 \left[\log(\sec x) \frac{x^2}{2} - \int \frac{1}{\sec x} \cdot \sec x \tan x \cdot \frac{x^2}{2} dx \right]$ $= x^2 \log(\sec x) - 2 \left[\log(\sec x) \frac{x^2}{2} - \frac{1}{2} \int x^2 \cdot \tan x \, dx \right]$ $= x^2 \log(\sec x) - 2 \left[\log(\sec x) \frac{x^2}{2} - \frac{1}{2} I \right]$ $I = x^2 \log(\sec x) - \log(\sec x) x^2 + I$ | <p>04</p> <p>½</p> <p>1</p> <p>½</p> <p>½</p> <p>½</p> |
| | d) | <p>-----</p> <p>Evaluate $\int \frac{\sec^2 x}{(\tan x)(\tan x + 1)} dx$</p> | <p>04</p> |

Note: If students attempted to solve the question give appropriate marks.



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|-----------|---|---|
| 4. | Ans | $\int \frac{\sec^2 x}{(\tan x)(\tan x + 1)} dx$ <p>Put $\tan x = t$ $\therefore \sec^2 x dx = dt$ $\therefore \int \frac{1}{(t)(t+1)} dt$ $\frac{1}{(t)(t+1)} = \frac{A}{t} + \frac{B}{t+1}$ $\therefore 1 = A(t+1) + B(t)$ \therefore Put $t = 0$, $A = 1$ Put $t = -1$, $B = -1$ $\therefore \frac{1}{(t)(t+1)} = \frac{1}{t} - \frac{1}{t+1}$ $\therefore \int \frac{1}{(t)(t+1)} dt = \int \left(\frac{1}{t} - \frac{1}{t+1} \right) dt$ $= \log(t) - \log(t+1) + c$ $= \log(\tan x) - \log(\tan x + 1) + c$</p> | <p>1</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> |
| | e) Ans | <p>Evaluate: $\int_0^{\frac{\pi}{2}} \frac{1}{1 + \sqrt{\tan x}} dx$</p> $\int_0^{\frac{\pi}{2}} \frac{1}{1 + \sqrt{\tan x}} dx$ $= \int_0^{\frac{\pi}{2}} \frac{1}{1 + \frac{\sqrt{\sin x}}{\sqrt{\cos x}}} dx$ <p>Let $I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx \dots \dots \dots (1)$</p> | <p>04</p> |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|---|---|
| 4. | e) | $I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos\left(\frac{\pi}{2}-x\right)}}{\sqrt{\cos\left(\frac{\pi}{2}-x\right)} + \sqrt{\sin\left(\frac{\pi}{2}-x\right)}} dx$ $I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx \dots\dots\dots (2)$ <p>Add (1) and (2)</p> $I+I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\cos x}}{\sqrt{\cos x} + \sqrt{\sin x}} dx + \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$ $2I = \int_0^{\frac{\pi}{2}} \frac{\sqrt{\sin x} + \sqrt{\cos x}}{\sqrt{\sin x} + \sqrt{\cos x}} dx$ $2I = \int_0^{\frac{\pi}{2}} dx$ $2I = [x]_0^{\frac{\pi}{2}}$ $2I = \frac{\pi}{2} - 0$ $\therefore I = \frac{\pi}{4}$ | <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1/2</p> |
| 5 | a) | <p>Solve any TWO of the following:</p> <p>Find area bounded by the curve $y = x^2$ and the line $y = x$</p> <p>We have $y = x^2$ and $y = x$</p> $\therefore x^2 - x = 0$ $\therefore x(x-1) = 0$ $\therefore x = 0 \text{ or } x = 1$ $\text{Area} = \int_a^b (y_1 - y_2) dx$ $= \int_0^1 (x^2 - x) dx$ | <p>12</p> <p>06</p> <p>1</p> <p>1</p> |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|---|------------------|
| 5. | a) | $= \left[\frac{x^3}{3} - \frac{x^2}{2} \right]_0^1$ $= \left[\frac{1^3}{3} - \frac{1^2}{2} - 0 \right]$ $= -\frac{1}{6}$ <p>$\therefore A = \frac{1}{6}$ or 0.167 (\because Area is always +ve)</p> | 1 1 1 1 |
| | b) | Attempt the following: | 06 |
| | i) | From the differential equation by eliminating the arbitrary constant if | 03 |
| | Ans | $y = A \cos x + B \sin x.$ $y = A \cos x + B \sin x.$ $\frac{dy}{dx} = -A \sin x + B \cos x$ $\frac{d^2y}{dx^2} = -A \cos x - B \sin x$ $= -(A \cos x + B \sin x)$ $= -y$ $\frac{d^2y}{dx^2} + y = 0$ | 1 1 1 |
| | ii) | Solve $(1+x^2)dy - x^2.ydx = 0$ | 03 |
| | Ans | $(1+x^2)dy - x^2.ydx = 0$ $(1+x^2)dy = x^2.ydx$ $\frac{dy}{y} = \frac{x^2 dx}{1+x^2}$ $\int \frac{dy}{y} = \int \frac{x^2 dx}{1+x^2}$ $\int \frac{dy}{y} = \int \frac{1+x^2-1 dx}{1+x^2}$ | 1 |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code:

22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|-----------|--|-----------------------------|
| 5. | | $\int \frac{dy}{y} = \int \left[1 - \frac{1}{1+x^2} \right] dx$ $\log y = x - \tan^{-1} x + c$ | 1 1 |
| | c) Ans | <p>Solve the D.E $\frac{dq}{dt} + \frac{1}{RC}q = \frac{E}{R}$ given that $q = 0$ when $t = 0$ and E,R,C are constant</p> $\frac{dq}{dt} + \frac{1}{RC}q = \frac{E}{R}$ $I.F = e^{\int \frac{1}{RC} dt}$ $= e^{\frac{t}{RC}}$ $\therefore q.e^{\frac{t}{RC}} = \int \frac{E}{R}.e^{\frac{t}{RC}} dt$ $= \frac{E}{R} e^{\frac{t}{RC}} \cdot \frac{1}{\frac{1}{RC}} + c_1$ $q.e^{\frac{t}{RC}} = e^{\frac{t}{RC}} EC + c_1$ <p>given that $q = 0$ when $t = 0$</p> $0 = e^0 EC + c_1$ $c_1 = -EC$ $q.e^{\frac{t}{RC}} = e^{\frac{t}{RC}} EC - EC$ $q = EC \left(1 - e^{-\frac{t}{RC}} \right)$ | 06 1 1 1 1 1 |
| 6. | | <p>Solve any TWO of the following:</p> | 12 |
| | a) i) | <p>Attempt the following:</p> <p>Solve the equations by Gauss-Seidal method. (two iterations only)</p> $10x + y + 2z = 13, \quad 3x + 10y + z = 14, \quad 2x + 3y + 10z = 15$ | 06 03 |



WINTER- 19 EXAMINATION

Subject Name: Applied Mathematics

Model Answer

Subject Code: 22224

| Q. No. | Sub Q.N. | Answers | Marking Scheme |
|--------|----------|---|--|
| 6. | c) | <p>Initial root $x_0=2$ $\therefore f'(2)=5$ $x_1 = x_0 - \frac{f(x_0)}{f'(x_0)} = 2 - \frac{f(2)}{f'(2)} = 1.8$ $x_2 = 1.8 - \frac{f(1.8)}{f'(1.8)} = 1.7913$ $x_3 = 1.7913 - \frac{f(1.7913)}{f'(1.7913)} = 1.7912$ $x_4 = 1.7912 - \frac{f(1.7912)}{f'(1.7912)} = 1.7912$ OR Let $f(x) = x^2 + x - 5$ $f(1) = -3 < 0$ $f(2) = 1 > 0$ $f'(x) = 2x + 1$ Initial root $x_0=2$ $x_i = x - \frac{f(x)}{f'(x)} = x - \frac{x^2 + x - 5}{2x + 1}$ $= \frac{2x^2 + x - x^2 - x + 5}{2x + 1}$ $= \frac{x^2 + 5}{2x + 1}$ $x_1 = 1.8$ $x_2 = 1.7913$ $x_3 = 1.7912$ $x_4 = 1.7912$</p> | <p>1 1 1 1 1 1 2 ½ ½ ½ ½</p> |
| | | <p><u>Important Note</u></p> <p><i>In the solution of the question paper, wherever possible all the possible alternative methods of solution are given for the sake of convenience. Still student may follow a method other than the given herein. In such case, first see whether the method falls within the scope of the curriculum, and then only give appropriate marks in accordance with the scheme of marking.</i></p> | |